

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C12N 15/10, A01H 5/00, C12N 15/53, C12N 15/82	A3	(11) International Publication Number: WO 00/20573 (43) International Publication Date: 13 April 2000 (13.04.2000)
(21) International Application Number: PCT/US99/23385 (22) International Filing Date: 06 October 1999 (06.10.1999) (30) Priority Data: 60/103,441 07 October 1998 (07.10.1998) US (60) Parent Application or Grant MAXYGEN, INC. [/]; O. SUBRAMANIAN, Venkiteswaran [/]; O. SUBRAMANIAN, Venkiteswaran [/]; O. QUINE, Jonathan, Alan ; O.		Published
(54) Title: DNA SHUFFLING TO PRODUCE NUCLEIC ACIDS FOR MYCOTOXIN DETOXIFICATION (54) Titre: REARRANGEMENT D'ADN POUR PRODUIRE DES ACIDES NUCLEIQUES DE DETOXICATION CONTRE LES MYCOTOXINES		
(57) Abstract Methods of shuffling nucleic acids to acquire or enhance mycotoxin detoxification activity, libraries of shuffled mycotoxin detoxification nucleic acids, transgenic cells, plants and DNA shuffling mixtures are provided.		
(57) Abrégé La présente invention concerne des procédés de réarrangement d'acides nucléiques permettant d'obtenir ou d'augmenter l'activité de détoxification contre les mycotoxines, des bibliothèques d'acide nucléiques de détoxification contre les mycotoxines, des cellules transgéniques, ainsi que des mélanges de réarrangement d'ADN et de plantes.		

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/10, 15/53, 15/82, A01H 5/00		(11) International Publication Number: WO 00/20573
A3		(43) International Publication Date: 13 April 2000 (13.04.00)
(21) International Application Number: PCT/US99/23385 (22) International Filing Date: 6 October 1999 (06.10.99) (30) Priority Data: 60/103,441 7 October 1998 (07.10.98) US (71) Applicant (for all designated States except US): MAXYGEN, INC. [US/US]; 515 Galveston Drive, Redwood City, CA 94063 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): SUBRAMANIAN, Venkiteswaran [US/US]; 3980 Corte Mar de Hierba, San Diego, CA 92130 (US). (74) Agents: QUINE, Jonathan, Alan; The Law Offices of Jonathan Alan Quine, P.O. Box 458, Alameda, CA 94501 (US) et al.		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. (88) Date of publication of the international search report: 20 July 2000 (20.07.00)
(54) Title: DNA SHUFFLING TO PRODUCE NUCLEIC ACIDS FOR MYCOTOXIN DETOXIFICATION		
(57) Abstract Methods of shuffling nucleic acids to acquire or enhance mycotoxin detoxification activity, libraries of shuffled mycotoxin detoxification nucleic acids, transgenic cells, plants and DNA shuffling mixtures are provided.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/23385A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/10 C12N15/53 C12N15/82 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01H

Documentation searched other than minimum: documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 23758 A (INCYTE PHARMA INC ;GOLI SURYA K (US); HILLMAN JENNIFER L (US)) 4 June 1998 (1998-06-04) page 2, line 7 - line 12 page 2, line 24 - line 29 page 16, line 28 -page 17, line 6 page 25, line 6 - line 9	1-21, 27-32
X	WO 96 06175 A (PIONEER HI BRED INT) 29 February 1996 (1996-02-29) the whole document --- -/--	9,22-26

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

5 April 2000

Date of mailing of the international search report

11/04/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 sport,
Fax: (+31-70) 340-3016

Authorized officer

Maddox, A

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/23385

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DUVICK, J., ET AL.: "Detoxification of mycotoxins in plants as a strategy for improving grain quality and disease resistance: Identification of fumonisin-degrading microbes from maize" MOLECULAR GENETICS OF HOST-SPECIFIC TOXINS IN PLANT DISEASE. PROCEEDINGS OF THE 3RD TOTTORI INTERNATIONAL SYMPOSIUM DAISEN, TOTTORI, JAPAN 24-29 AUGUST 1997(1998). KLUWER ACADEMIC PRESS, DORDRECHT, pages 369-381, XP002121275 the whole document</p>	1-32
A	<p>WO 96 12414 A (ERBER ERICH KG) 2 May 1996 (1996-05-02) the whole document</p>	1-32
A	<p>KIMURA M ET AL: "Trichothecene 3-O-acetyltransferase protects both the producing organism and transformed yeast from related mycotoxins. Cloning and characterization of Tr1101" JOURNAL OF BIOLOGICAL CHEMISTRY, US, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, vol. 273, no. 3, 16 January 1998 (1998-01-16), pages 1654-1661-1661, XP002125091 ISSN: 0021-9258 the whole document</p>	1-32
A	<p>WO 97 35966 A (MAXYGEN INC ;MINSHULL JEREMY (US); STEMMER WILLEM P C (US)) 2 October 1997 (1997-10-02) the whole document</p>	1-32
A	<p>WO 97 20078 A (AFFYMAX TECH NV ;CRAMERI ANDREAS (US); STEMMER WILLEM P C (US)) 5 June 1997 (1997-06-05) the whole document</p>	1-32
A	<p>HARAYAMA S: "Artificial evolution by DNA shuffling" TRENDS IN BIOTECHNOLOGY, GB, ELSEVIER PUBLICATIONS, CAMBRIDGE, vol. 16, no. 2, February 1998 (1998-02), pages 76-82, XP004107046 ISSN: 0167-7799 the whole document</p>	1-32

-/--

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/23385

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO 00 04158 A (DUVICK JONATHAN P ;PIONEER HI BRED INT (US); GILLIAM JACOB (US); M) 27 January 2000 (2000-01-27) page 12, line 21 -page 13, line 5 -----	1-4,9
E	WO 00 04160 A (PIONEER HI BRED INT ;FOLKERTS OTTO (US); CRASTA OSWALD R (US); CUR) 27 January 2000 (2000-01-27) page 26, line 25 -page 27, line 18 -----	1-4,9

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/23385

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9823758 A	04-06-1998	US 5817497 A	06-10-1998
		AU 5181198 A	22-06-1998
		US 5976528 A	02-11-1999
WO 9606175 A	29-02-1996	US 5792931 A	11-08-1998
		AU 697831 B	15-10-1998
		AU 3491095 A	14-03-1996
		AU 9725398 A	15-04-1999
		CA 2195784 A	29-02-1996
		EP 0775211 A	28-05-1997
		EP 0981953 A	01-03-2000
		NZ 292668 A	25-11-1998
		US 5716820 A	10-02-1998
		US 6025188 A	15-02-2000
		ZA 9506531 A	25-03-1996
WO 9612414 A	02-05-1996	AT 504 U	27-12-1995
		AT 175073 T	15-01-1999
		AU 3736595 A	15-05-1996
		BR 9509399 A	04-11-1997
		CA 2192983 A	02-05-1996
		CN 1158554 A	03-09-1997
		CZ 9603683 A	11-06-1997
		DE 59504703 D	11-02-1999
		EP 0786945 A	06-08-1997
		ES 2128086 T	01-05-1999
		HR 950526 A	30-04-1997
		HU 76374 A	28-08-1997
		IL 115650 A	14-07-1999
		JP 2983639 B	29-11-1999
		JP 9511657 T	25-11-1997
		PL 319755 A	18-08-1997
		SK 167896 A	04-06-1997
		ZA 9508786 A	13-05-1996
WO 9735966 A	02-10-1997	US 5837458 A	17-11-1998
		AU 2337797 A	17-10-1997
		AU 2542697 A	17-10-1997
		CA 2247930 A	02-10-1997
		EP 0932670 A	04-08-1999
		EP 0906418 A	07-04-1999
		WO 9735957 A	02-10-1997
		AU 713952 B	16-12-1999
		AU 1087397 A	19-06-1997
		CA 2239099 A	05-06-1997
		EP 0876509 A	11-11-1998
		EP 0911396 A	28-04-1999
		JP 2000500981 T	02-02-2000
WO 9720078 A	05-06-1997	WO 9720078 A	05-06-1997
		US 5811238 A	22-09-1998
		AU 713952 B	16-12-1999
		AU 1087397 A	19-06-1997
		AU 2542697 A	17-10-1997
		CA 2239099 A	05-06-1997
		EP 0876509 A	11-11-1998
		EP 0906418 A	07-04-1999
		EP 0911396 A	28-04-1999

Form PCT/ISA/210 (patent family annex) (July 1992)

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 99/23385

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9720078 A		JP 2000500981 T	02-02-2000
		WO 9735966 A	02-10-1997
		US 5837458 A	17-11-1998
WO 9831816 A	23-07-1998	AU 6029998 A	07-08-1998
WO 9831837 A	23-07-1998	AU 5920998 A	07-08-1998
WO 9827230 A	25-06-1998	AU 5729298 A	15-07-1998
		EP 0946755 A	06-10-1999
WO 0004158 A	27-01-2000	NONE	
WO 0004160 A	27-01-2000	WO 0004159 A	27-01-2000



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C12N 15/10, A01H 5/00, C12N 15/53, C12N 15/82	A2	(11) International Publication Number: WO 00/20573 (43) International Publication Date: 13 April 2000 (13.04.2000)
(21) International Application Number: PCT/US99/23385 (22) International Filing Date: 06 October 1999 (06.10.1999) (30) Priority Data: 60/103,441 07 October 1998 (07.10.1998) US (60) Parent Application or Grant MAXYGEN, INC. [/]; (). SUBRAMANIAN, Venkiteswaran [/]; (). SUBRAMANIAN, Venkiteswaran [/]; (). QUINE, Jonathan, Alan ; ().		Published
(54) Title: DNA SHUFFLING TO PRODUCE NUCLEIC ACIDS FOR MYCOTOXIN DETOXIFICATION (54) Titre: REARRANGEMENT D'ADN POUR PRODUIRE DES ACIDES NUCLEIQUES DE DETOXICATION CONTRE LES MYCOTOXINES (57) Abstract <p>Methods of shuffling nucleic acids to acquire or enhance mycotoxin detoxification activity, libraries of shuffled mycotoxin detoxification nucleic acids, transgenic cells, plants and DNA shuffling mixtures are provided.</p> (57) Abrégé <p>La présente invention concerne des procédés de réarrangement d'acides nucléiques permettant d'obtenir ou d'augmenter l'activité de détoxification contre les mycotoxines, des bibliothèques d'acide nucléiques de détoxification contre les mycotoxines, des cellules transgéniques, ainsi que des mélanges de réarrangement d'ADN et de plantes.</p>		

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/10, 15/53, 15/82, A01H 5/00	A2	(11) International Publication Number: WO 00/20573 (43) International Publication Date: 13 April 2000 (13.04.00)
(21) International Application Number: PCT/US99/23385 (22) International Filing Date: 6 October 1999 (06.10.99) (30) Priority Data: 60/103,441 7 October 1998 (07.10.98) US (71) Applicant (for all designated States except US): MAXYGEN, INC. [US/US]; 515 Galveston Drive, Redwood City, CA 94063 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): SUBRAMANIAN, Venkiteswaran [US/US]; 3980 Corte Mar de Hierba, San Diego, CA 92130 (US). (74) Agents: QUINE, Jonathan, Alan; The Law Offices of Jonathan Alan Quine, P.O. Box 458, Alameda, CA 94501 (US) et al.		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: DNA SHUFFLING TO PRODUCE NUCLEIC ACIDS FOR MYCOTOXIN DETOXIFICATION (57) Abstract Methods of shuffling nucleic acids to acquire or enhance mycotoxin detoxification activity, libraries of shuffled mycotoxin detoxification nucleic acids, transgenic cells, plants and DNA shuffling mixtures are provided.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	IS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CC	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

4

5

Description

5

10

15

20

25

30

35

40

45

50

55

DNA SHUFFLING TO PRODUCE NUCLEIC ACIDS FOR MYCOTOXIN DETOXIFICATION

CROSS REFERENCE TO RELATED APPLICATIONS

The present application is a non-provisional of "DNA SHUFFLING TO PRODUCE NUCLEIC ACIDS FOR MYCOTOXIN DETOXIFICATION" by Subramanian, USSN 60/103,441, filed October 7, 1998.

FIELD OF THE INVENTION

This invention pertains to the shuffling of nucleic acids to achieve or enhance mycotoxin detoxification, especially in plants.

BACKGROUND OF THE INVENTION

"Mycotoxins" generically refer to a number of toxic molecules produced by fungal species, such as polyketides (including aflatoxins, demethylsterigmatocystin, O-methylsterigmatocystin etc.), fumonisins, alperisins (e.g., A₁, A₂, B₁, B₂), sphingofungins (A, B, C and D), trichothecenes, fumifungins, and the like. Polyketides are a large structurally diverse class of secondary metabolites synthesized by bacteria, fungi, and plants and are formed by a polyketide synthase (PKS) through the sequential condensation of small carboxylic acids. Katz and Donadio (1993) Annu Rev. Microbiol. 47:875-912; Brown et al. (1996) PNAS 93:14873-14877; Silva et al. (1996) J. Biol Chem. 271: 13600-608.

Aflatoxin B₁ is the principal member of the aflatoxin (AF) family of polyketide mycotoxins produced by *Aspergillus parasiticus*, *Aspergillus flavus* and *Aspergillus nomius*. Aflatoxin B₁ is the most potent mycotoxin known to man. For example, AF was characterized as the causative agent for the death of more than a hundred thousand poultry in England that had ingested AF-contaminated peanut meal. This discovery led to legislation regulating the trade of AF-contaminated agricultural commodities.

Sterigmatocystin (ST) is a related polyketide mycotoxin, which is produced by several members of the *Aspergillus*. ST is the second to last intermediate in

the biosynthesis of AF. Kelkar et al. (1997) J. Biol Chem. 272: 1589-94. Various *Aspergillus* species that produce AF and ST are known to be pathogenic to corn, grains and nuts and are known to produce these mycotoxins during the growth of the crops and during storage, leading to the introduction of AF and ST into primary food stuffs. AF and ST are acutely toxic and carcinogenic and are a serious concern from human and animal health perspective. Busby & Wogan (1985) in Chemical Carcinogens (Searle ed., 1985) pp 945-1136, American Chemical Society, Washington D.C.

Trichothecenes are another family of sesquiterpenoid mycotoxins produced by *Fusarium* species and other molds that are known plant pathogens. These compounds are potent inhibitors of protein synthesis in eukaryotes (Kimura et al. (1998) J. Biol Chem. 273: 1654-1661) and reportedly bind to the 60S ribosomal subunits to prevent polypeptide chain initiation or elongation. Trichothecenes are also an important group of mycotoxins that cause serious problems of food pollution. They have been implicated in incidents of mycotoxicosis including vomiting, dermatitis and hemorrhagic septicemia in humans and livestock, resulting in loss of productivity and even death. Lastly, fumonisins (F) are another structurally distinct class of mycotoxins produced by several *Fusarium* species that is involved in food poisoning and toxic effects. Scott (1993) International Journal of Food Microbiology 18:257-270 and the references therein provide a review of the Fumonisins.

Thus, the contamination of corn, grains and nuts with various types of mycotoxins produced by pathogenic species such as *Aspergillus* and *Fusarium* is a major health and food pollution problem, as well as causing reduction in crop yields by being toxic to infected plants. These mycotoxins survive food processing, which adds to the problem. It is well known that ST and AF induce liver cancer and are linked to a specific mutation in a tumor suppressor gene. Brown et al. (1996) PNAS 93: 14873-14877. Natural aflatoxins and other mycotoxins like ST do not pose a major health threat *per se*; however, renal and hepatic oxidative detoxification of these compounds in contaminated foods by cytochrome P450 enzymes yields an epoxide which is cytotoxic.

For example, AFB1 is converted to its 15,16-exo-epoxide, which is a highly toxic mutagen. Silva et al. (1996), *supra* and references therein. It has been shown that this epoxide targets guanine residues and selectively alkylates the N-7 position

5 of this purine in double-stranded DNA. Depurination of the alkylated base has been
correlated to bladder cancer in laboratory mice, teratogenic effects in chicken embryos
and liver cancer in humans. A direct correlation between DNA damage and human
10 cancer has been established and is related to the mutational hot spots of p53, an important
5 tumor suppressor gene. Approximately 50% of all cancers have associated altered p53
sequences.

Trading of AF-contaminated agricultural commodities is tightly regulated
15 at both national and international levels. Compliance to these regulations causes the loss
of millions of dollars in agricultural produce in US each year. Trade sanctions and
10 health effects on mycotoxin contaminated grains add significantly to the losses (Brown et
al. (1996) PNAS 93: 14873-14877).

Accordingly, it is highly desirable to transform various mycotoxins
produced by fungal pathogens in various crops into inactive compounds with respect to
plant, human and animal toxicity. This would alleviate important food pollution
25 problems, as well as cost associated with complying with detecting AF-contamination in
various crop commodities and destroying them. Surprisingly, the present invention
provides for the detoxification of mycotoxis by transformation of the mycotoxins into
30 non-toxic compounds. This detoxification is particularly useful in crops, thereby solving
each of the problems outlined above, as well as providing a variety of other features
20 which will be apparent upon review.

SUMMARY OF THE INVENTION

35 In the present invention, DNA shuffling is used to generate new or
improved mycotoxin detoxification genes. These mycotoxin detoxification genes are
used to provide enzymes which degrade mycotoxins, in agricultural and industrial
40 25 processes. These new and/or improved genes have surprisingly superior properties as
compared to naturally occurring mycotoxin detoxification genes.

In the methods for obtaining mycotoxin resistant genes, a plurality of
parental forms (homologs) of a selected nucleic acid are recombined. The selected
45 nucleic acid is derived either from one or more parental nucleic acid(s) which encodes an
enzyme which degrades or modifies a mycotoxin, or a fragment thereof, or from a
30 parental nucleic acid which does not encode mycotoxin detoxification, but which is a

5 substrate for DNA shuffling to develop monooxygenase activity. The plurality of forms
of the selected nucleic acid differ from each other in at least one (and typically two or
10 more) nucleotides, and, upon recombination, provide a library of recombinant mycotoxin
detoxification nucleic acids. The library can be an *in vitro* set of molecules, or present in
5 cells, phage or the like. The library is screened to identify at least one recombinant
mycotoxin detoxification nucleic acid that exhibits distinct or improved mycotoxin
detoxification activity (typically in an encoded polypeptide) compared to the parental
15 nucleic acid or nucleic acids.

In selecting for mycotoxin detoxification activity, a candidate shuffled
10 DNA can be tested for encoded mycotoxin detoxification activity in essentially any
process. Common processes that can be screened include screening for inactivation or
20 modification of an aflatoxin, inactivation or modification of a sterigmatocystin,
inactivation or modification of a trichothecene, and inactivation or modification of a
fumonisin. Similarly, instead of, or in addition to, testing for an increase in mycotoxin
25 detoxification activity, it is also desirable to screen for shuffled nucleic acids which
produce higher levels of a mycotoxin detoxification nucleic acid or enhanced or reduced
recombinant mycotoxin detoxification polypeptide expression, or increased stability
30 encoded by the recombinant mycotoxin resistant nucleic acid.

A variety of screening methods can be used to screen a library, depending
20 on the mycotoxin detoxification activity for which the library is selected. By way of
example, the library to be screened can be present in a population of cells. The library is
35 selected by growing the cells in or on a medium comprising the mycotoxin to be
degraded and selecting for a detected physical difference between, e.g., oxidized or
reduced forms of the mycotoxin and the non-oxidized or reduced form of the mycotoxin,
40 25 either in the cell, or the extracellular medium. Alternately, survival of library cells on a
medium which includes a mycotoxin can be used to screen the library.

Iterative selection for mycotoxin detoxification nucleic acids is also a
45 feature of the invention. In these methods, a selected nucleic acid identified as encoding
mycotoxin detoxification activity can be shuffled, either with the parental nucleic acids,
30 or with other nucleic acids (e.g., mutated forms of the selected nucleic acid) to produce a
second shuffled library. The second shuffled library is then selected for one or more
50

5 form of mycotoxin detoxification activity, which can be the same or different than the mycotoxin detoxification activity previously selected.

10 This process can be iteratively repeated as many times as desired, until a nucleic acid with optimized or desired mycotoxin detoxification properties is obtained. If
5 desired, any nucleic acid identified by any of the methods herein can be cloned and, optionally, expressed. Because of the need to reduce mycotoxin pollution/contamination of foods, it is desirable to express mycotoxin detoxification nucleic acids in, e.g., plants,
15 thereby reducing the occurrence of mycotoxins in the plants. Furthermore, mycotoxin detoxification in plants also adds to the vigor of the plants.

10 The invention also provides methods of increasing mycotoxin detoxification activity by whole genome shuffling. In these methods, a plurality of genomic nucleic acids are shuffled in a cell (in whole cell shuffling, entire genomes are shuffled, rather than specific sequences, although "spiking" of selected nucleic acids can
20 be used to bias shuffling outcomes). The resulting shuffled nucleic acids are selected for one or more mycotoxin detoxification traits. The genomic nucleic acids can be from a species or strain different from the cell in which activity is desired. Similarly, the
25 shuffling reaction can be performed in cells using genomic DNA from the same or different species, or strains. Strains or enzymes exhibiting enhanced activity can be identified.

20 The distinct or improved activity encoded by a nucleic acid identified after shuffling can encode one or more of a variety of properties, including, e.g., inactivation
35 or modification of a polyketide, an aflatoxin, inactivation or modification of a sterigmatocystin, inactivation or modification of a trichothecene, inactivation or modification of a fumonisin, an increased ability to chemically modify a mycotoxin, an
40 increase in the range of mycotoxin substrates which the distinct or improved nucleic acid operates on, an increased expression level of a polypeptide encoded by the nucleic acid, a decrease in susceptibility of a polypeptide encoded by the nucleic acid to protease
45 cleavage, a decrease in susceptibility of a polypeptide encoded by the nucleic acid to high or low pH levels, a decrease in susceptibility of the protein encoded by the nucleic acid to high or low temperatures, and a decrease in toxicity to a host cell of a polypeptide
30 encoded by the selected nucleic acid.

5 The selected nucleic acids to be shuffled can be from any of a variety of
sources, including synthetic or cloned DNAs. Exemplar targets for recombination
include: nucleic acids encoding a monooxygenase, a P450, trichothecene-3-O-
10 acetyltransferase, a 3-O-Methyltransferase, a glutathione S-transferase, an epoxide
5 hydrolase, an isomerase, a macrolide-O-acyltransferase, a 3-O-acyltransferase, and a
cis-diol producing monooxygenase which is specific for furan. Typically, shuffled
nucleic acids are cloned into expression vectors to achieve desired expression levels.

15 One feature of the invention is the production of libraries and shuffling
mixtures for use in the methods as set forth above. For example, a phage display library
10 comprising shuffled forms of a nucleic acid is provided. Similarly, a shuffling mixture
comprising at least three homologous DNAs, each of which is derived from a nucleic
20 acid encoding a polypeptide or polypeptide fragment, is provided. These polypeptides
can be, for example, any of those noted herein.

25 Isolated nucleic acids identified by selection of the libraries in the methods
15 above are also a feature of the invention, as are kits comprising any of: mycotoxin
detoxification nucleic acid libraries, shuffled mycotoxin detoxification nucleic acids,
instructional materials for practicing any of the methods herein, containers for holding
30 other kit components, and the like.

BRIEF DESCRIPTION OF THE FIGURES

20 Not Applicable.

DEFINITIONS

35 Unless clearly indicated to the contrary, the following definitions
supplement definitions of terms known in the art.

40 A "recombinant monooxygenase nucleic acid" is a recombinant nucleic
25 acid encoding a protein or RNA which confers monooxygenase activity to a cell when the
nucleic acid is expressed in the cell.

45 A "recombinant" nucleic acid is a nucleic acid produced by recombination
between two or more nucleic acids, or any nucleic acid made by an *in vitro* or artificial
process. The term "recombinant" when used with reference to a cell indicates that the
30 cell comprises (and optionally replicates) a heterologous nucleic acid, or expresses a
peptide or protein encoded by a heterologous nucleic acid. Recombinant cells can

5 contain genes that are not found within the native (non-recombinant) form of the cell.
Recombinant cells can also contain genes found in the native form of the cell where the
genes are modified and re-introduced into the cell by artificial means. The term also
10 encompasses cells that contain a nucleic acid endogenous to the cell that has been
5 artificially modified without removing the nucleic acid from the cell; such modifications
include those obtained by gene replacement, site-specific mutation, and related
15 techniques.

A "recombinant mycotoxin detoxification nucleic acid" is a recombinant
nucleic acid encoding a protein or RNA which confers mycotoxin detoxification or
10 degradation activity to a cell when the nucleic acid is expressed in the cell (and, most
20 typically, translated into a polypeptide).

A "plurality of forms" of a selected nucleic acid refers to a plurality of
homologs of the nucleic acid. The homologs can be from naturally occurring homologs
25 (e.g., two or more homologous genes) or by artificial synthesis of one or more nucleic
15 acids having related sequences, or by modification of one or more nucleic acid to
produce related nucleic acids. Nucleic acids are homologous when they are derived,
naturally or artificially, from a common ancestor sequence. During natural evolution,
30 this occurs when two or more descendent sequences diverge from a parent sequence over
time, i.e., due to mutation and natural selection. Under artificial conditions, divergence
20 occurs, e.g., in one of two ways. First, a given sequence can be artificially recombined
with another sequence, as occurs, e.g., during typical cloning, to produce a descendent
35 nucleic acid. Alternatively, a nucleic acid can be synthesized *de novo*, by synthesizing a
nucleic acid which varies in sequence from a given parental nucleic acid sequence.

When there is no explicit knowledge about the ancestry of two nucleic
40 25 acids, homology is typically inferred by sequence comparison between two sequences.
Where two nucleic acid sequences show sequence similarity it is inferred that the two
nucleic acids share a common ancestor. The precise level of sequence similarity required
45 to establish homology varies in the art depending on a variety of factors. For purposes of
this disclosure, two sequences are considered homologous where they share sufficient
30 sequence identity to allow direct recombination to occur between two nucleic acid
molecules (as opposed to recombination using oligonucleotide intermediates, which does

5 not require sequence similarity to achieve recombination). Typically, nucleic acids
require regions of close similarity spaced roughly the same distance apart to permit
10 recombination to occur. The recombination can be in vitro or in vivo.

10 The terms "identical" or percent "identity," in the context of two or more
5 nucleic acid or polypeptide sequences, refer to two or more sequences or subsequences
that are the same or have a specified percentage of amino acid residues or nucleotides
that are the same, when compared and aligned for maximum correspondence, as
15 measured using one of the sequence comparison algorithms described below (or other
algorithms available to persons of skill) or by visual inspection.

10 The phrase "substantially identical," in the context of two nucleic acids or
20 polypeptides (e.g., DNAs encoding a monooxygenase, or the amino acid sequence of the
monooxygenase) refers to two or more sequences or subsequences that have at least about
60%, preferably 80%, most preferably 90-95% nucleotide or amino acid residue identity,
when compared and aligned for maximum correspondence, as measured using one of the
25 following sequence comparison algorithms or by visual inspection. Such "substantially
15 identical" sequences are typically considered to be homologous. Preferably, the
"substantial identity" exists over a region of the sequences that is at least about 50
30 residues in length, more preferably over a region of at least about 100 residues, and most
preferably the sequences are substantially identical over at least about 150 residues, or
20 over the full length of the two sequences to be compared.

35 For sequence comparison and homology determination, typically one
sequence acts as a reference sequence to which test sequences are compared. When
using a sequence comparison algorithm, test and reference sequences are input into a
computer, subsequence coordinates are designated, if necessary, and sequence algorithm
40 25 program parameters are designated. The sequence comparison algorithm then calculates
the percent sequence identity for the test sequence(s) relative to the reference sequence,
based on the designated program parameters.

45 Optimal alignment of sequences for comparison can be conducted, e.g., by
the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by
30 the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443
(1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad.*

5 *Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by visual inspection (*see generally*,
10 Ausubel *et al.*, *infra*).

5 One example of algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul *et al.*, *J. Mol. Biol.* 215:403-410 (1990). Software for performing BLAST analyses is
15 publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring
10 sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a
20 word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, *supra*). These initial neighborhood word hits act as
25 seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment
15 score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0)
30 and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in
20 each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the
35 accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the
sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences)
40 uses as defaults a wordlength (W) of 11, an expectation (E) of 10, a cutoff of 100, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP
25 program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (*see* Henikoff & Henikoff (1989) *Proc. Natl. Acad. Sci. USA*
45 89:10915).

30 In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (*see, e.g.*,

5 Karlin & Altschul (1993) *Proc. Nat'l. Acad. Sci. USA* 90:5873-5787). One measure of
similarity provided by the BLAST algorithm is the smallest sum probability (P(N)),
which provides an indication of the probability by which a match between two nucleotide
10 or amino acid sequences would occur by chance. For example, a nucleic acid is
5 considered similar to a reference sequence if the smallest sum probability in a comparison
of the test nucleic acid to the reference nucleic acid is less than about 0.1, more
preferably less than about 0.01, and most preferably less than about 0.001.

15 Another indication that two nucleic acid sequences are substantially
identical/ homologous is that the two molecules hybridize to each other under stringent
10 conditions. The phrase "hybridizing specifically to," refers to the binding, duplexing, or
20 hybridizing of a molecule only to a particular nucleotide sequence under stringent
conditions, including when that sequence is present in a complex mixture (*e.g.*, total
cellular) DNA or RNA. "Bind(s) substantially" refers to complementary hybridization
25 between a probe nucleic acid and a target nucleic acid and embraces minor mismatches
15 that can be accommodated by reducing the stringency of the hybridization media to
achieve the desired detection of the target polynucleotide sequence.

30 "Stringent hybridization conditions" and "stringent hybridization wash
conditions" in the context of nucleic acid hybridization experiments such as Southern and
northern hybridizations are sequence dependent, and are different under different
20 environmental parameters. Longer sequences hybridize specifically at higher
temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen
35 (1993) *Laboratory Techniques in Biochemistry and Molecular Biology--Hybridization
with Nucleic Acid Probes* part I chapter 2 "Overview of principles of hybridization and
the strategy of nucleic acid probe assays," Elsevier, New York. Generally, highly
40 25 stringent hybridization and wash conditions are selected to be about 5° C lower than the
thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH.
Typically, under "stringent conditions" a probe will hybridize to its target subsequence,
45 but no to unrelated sequences.

50 The T_m is the temperature (under defined ionic strength and pH) at which
30 50% of the target sequence hybridizes to a perfectly matched probe. Very stringent
conditions are selected to be equal to the T_m for a particular probe. An example of

5 stringent hybridization conditions for hybridization of complementary nucleic acids which
have more than 100 complementary residues on a filter in a Southern or northern blot is
10 50% formamide with 1 mg of heparin at 42°C, with the hybridization being carried out
overnight. An example of highly stringent wash conditions is 0.15M NaCl at 72°C for
10 about 15 minutes. An example of stringent wash conditions is a 0.2x SSC wash at 65°C
5 for 15 minutes (*see*, Sambrook, *infra.*, for a description of SSC buffer). Often, a high
stringency wash is preceded by a low stringency wash to remove background probe
15 signal. An example medium stringency wash for a duplex of, *e.g.*, more than 100
nucleotides, is 1x SSC at 45°C for 15 minutes. An example low stringency wash for a
10 duplex of, *e.g.*, more than 100 nucleotides, is 4-6x SSC at 40°C for 15 minutes. For
short probes (*e.g.*, about 10 to 50 nucleotides), stringent conditions typically involve salt
20 concentrations of less than about 1.0 M Na ion, typically about 0.01 to 1.0 M Na ion
concentration (or other salts) at pH 7.0 to 8.3, and the temperature is typically at least
about 30°C. Stringent conditions can also be achieved with the addition of destabilizing
25 agents such as formamide. In general, a signal to noise ratio of 2x (or higher) than that
observed for an unrelated probe in the particular hybridization assay indicates detection
of a specific hybridization.

30 A further indication that two nucleic acid sequences or polypeptides are
substantially identical/homologous is that the polypeptide encoded by the first nucleic
20 acid is immunologically cross reactive with, or specifically binds to, the polypeptide
encoded by the second nucleic acid. Thus, a polypeptide is typically substantially
35 identical to a second polypeptide, for example, where the two peptides differ only by
conservative substitutions.

40 "Conservatively modified variations" of a particular polynucleotide
25 sequence refers to those polynucleotides that encode identical or essentially identical
amino acid sequences, or where the polynucleotide does not encode an amino acid
sequence, to essentially identical sequences. Because of the degeneracy of the genetic
45 code, a large number of functionally identical nucleic acids encode any given
polypeptide. For instance, the codons CGU, CGC, CGA, CGG, AGA, and AGG all
30 encode the amino acid arginine. Thus, at every position where an arginine is specified by
a codon, the codon can be altered to any of the corresponding codons described without

5 altering the encoded polypeptide. Such nucleic acid variations are "silent variations,"
which are one species of "conservatively modified variations." Every polynucleotide
sequence described herein which encodes a polypeptide also describes every possible
10 silent variation, except where otherwise noted. One of skill will recognize that each
5 codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine)
can be modified to yield a functionally identical molecule by standard techniques.
Accordingly, each "silent variation" of a nucleic acid which encodes a polypeptide is
15 implicit in each described sequence.

Furthermore, one of skill will recognize that individual substitutions,
10 deletions or additions which alter, add or delete a single amino acid or a small percentage
of amino acids (typically less than 5%, more typically less than 1%) in an encoded
20 sequence are "conservatively modified variations" where the alterations result in the
substitution of an amino acid with a chemically similar amino acid. Conservative
substitution tables providing functionally similar amino acids are well known in the art.
25 The following five groups each contain amino acids that are conservative substitutions for
one another: Aliphatic: Glycine (G), Alanine (A), Valine (V), Leucine (L), Isoleucine
(I); Aromatic: Phenylalanine (F), Tyrosine (Y), Tryptophan (W); Sulfur-containing:
30 Methionine (M), Cysteine (C); Basic: Arginine (R), Lysine (K), Histidine (H); Acidic:
Aspartic acid (D), Glutamic acid (E), Asparagine (N), Glutamine (Q). *See also*,
20 Creighton (1984) *Proteins*, W.H. Freeman and Company.

In addition, individual substitutions, deletions or additions which alter, add
35 or delete a single amino acid or a small percentage of amino acids in an encoded
sequence are also "conservatively modified variations." Sequences that differ by
conservative variations are generally homologous.

40 25 A "subsequence" refers to a sequence of nucleic acids or amino acids that
comprise a part of a longer sequence of nucleic acids or amino acids (*e.g.*, polypeptide)
respectively.

The term "gene" is used broadly to refer to any segment of DNA
45 associated with expression of a given RNA or protein. Thus, genes include regions
encoding expressed RNAs (which typically include polypeptide coding sequences) and,
30 often, the regulatory sequences required for their expression. Genes can be obtained

5 from a variety of sources, including cloning from a source of interest or synthesizing
from known or predicted sequence information, and may include sequences designed to
have desired parameters.

10 The term "isolated", when applied to a nucleic acid or protein, denotes
5 that the nucleic acid or protein is essentially free of other cellular components with which
it is associated in the natural state.

15 The term "nucleic acid" refers to deoxyribonucleotides or ribonucleotides
and polymers thereof in either single- or double-stranded form. Unless specifically
limited, the term encompasses nucleic acids containing known analogues of natural
10 nucleotides which have similar binding properties as the reference nucleic acid and are
metabolized in a manner similar to naturally occurring nucleotides. Unless otherwise
20 indicated, a particular nucleic acid sequence also implicitly encompasses conservatively
modified variants thereof (*e.g.* degenerate codon substitutions) and complementary
sequences and as well as the sequence explicitly indicated. Specifically, degenerate
25 codon substitutions may be achieved by generating sequences in which the third position
of one or more selected (or all) codons is substituted with mixed-base and/or
deoxyinosine residues (Batzer *et al.* (1991) *Nucleic Acid Res.* 19: 5081; Ohtsuka *et al.*
30 (1985) *J. Biol. Chem.* 260: 2605-2608; Cassol *et al.* (1992); Rossolini *et al.* (1994) *Mol.*
Cell. Probes 8: 91-98). The term nucleic acid is generic to the terms "gene", "DNA,"
20 "cDNA", "oligonucleotide," "RNA," "mRNA," and the like.

35 "Nucleic acid derived from a gene" refers to a nucleic acid for whose
synthesis the gene, or a subsequence thereof, has ultimately served as a template. Thus,
an mRNA, a cDNA reverse transcribed from an mRNA, an RNA transcribed from that
cDNA, a DNA amplified from the cDNA, an RNA transcribed from the amplified DNA,
40 25 *etc.*, are all derived from the gene and detection of such derived products is indicative of
the presence and/or abundance of the original gene and/or gene transcript in a sample.

45 A nucleic acid is "operably linked" when it is placed into a functional
relationship with another nucleic acid sequence. For instance, a promoter or enhancer is
operably linked to a coding sequence if it increases the transcription of the coding
30 sequence.

5 A "recombinant expression cassette" or simply an "expression cassette" is
a nucleic acid construct, generated recombinantly or synthetically, with nucleic acid
elements that are capable of effecting expression of a structural gene in hosts compatible
10 with such sequences. Expression cassettes include at least promoters and optionally,
5 transcription termination signals. Typically, the recombinant expression cassette includes
a nucleic acid to be transcribed (*e.g.*, a nucleic acid encoding a desired polypeptide), and
a promoter. Additional factors necessary or helpful in effecting expression may also be
15 used as described herein. For example, an expression cassette can also include
nucleotide sequences that encode a signal sequence that directs secretion of an expressed
10 protein from the host cell. Transcription termination signals, enhancers, and other
nucleic acid sequences that influence gene expression, can also be included in an
20 expression cassette.

DETAILED DISCUSSION OF THE INVENTION

25 It is highly desirable to enzymatically transform various mycotoxins
15 produced by fungal pathogens, into inactive compounds with respect to plant, human and
animal toxicity. This eliminates reductions in crop yield and subsequent food pollution
by mycotoxins, as well as costs associated with complying with detecting AF-
30 contamination in various crop commodities and destroying them.

Polyketides are synthesized in fungi, *e.g.*, by polyketide synthase. The
20 enzyme facilitates the reiterative condensation of simple carboxylic acids; typically,
acetyl-CoA as a starter unit and malonyl-CoA serves as an extender unit. The
35 biosynthetic pathway for AF and ST is, approximately, as follows: a hexanoate starter
unit is converted into an initial polyketide precursor (octaketide) which is converted into
norsolorinic acid, which is converted into averantin, which is converted into averufanin,
40 25 which is converted into averufin, which is converted into versiconal hemiacetal acetate,
which is converted into versicolorin B, which is converted into versicolorin A, which is
converted into demethylsterigmatocystin, which is converted into ST, which is converted
45 into O-methylsterigmatocystin, which is converted into AFB₁. *See*, Yu (1995) Journal of
Bacteriology 177(16):4792-4800 and the references cited therein, and Silva (1996) JBC
30 271(23):13600-13608 and the references cited therein.

5 The chemical structures of the important mycotoxin polyketides Aflatoxin B₁, and compounds in the biosynthetic pathway for Aflatoxin B₁, including sterigmatocystins, norisolornic acid, and a variety of other compounds can be found, e.g., in Silva et al. (1996) *JBC* 271:23:13600-13608 and the references cited therein.

10 The double bond at carbon atoms 15,16 of AFB₁ is very important with respect to toxicity (Silva et al., 1996, *id*). This double bond is also susceptible to oxidations such as hydroxylation, epoxidation etc. These are monooxygenase catalyzed reactions. A number of monooxygenases, including P450s (*see* Ortiz de Montellano (ed.) 1995, Cytochrome P450 Structure and Mechanism and Biochemistry, Second Edition Plenum Press (New York and London), monooxygenase from *P. oleovorans* (J. Biol. Chem., 248, 1725-1730, 1973; May L. Am. Chem. Soc., 98, 7856-7858) and other homologous non-heme iron-sulfur monooxygenases from *Rhodococcus*, *Mycobacterium*, *ocardia*, *Pseudomonas* and *Bacillus*; heme-dependent peroxidases, iron-sulfur monooxygenases and quinone-dependent monooxygenases are known and can be recombined in the methods herein to provide mycotoxin resistant nucleic acids. Many of these detoxification enzymes confer increased hydrophilicity to the mycotoxin, thereby facilitating excretion, e.g., in mammals.

30 P450s are particularly preferred monooxygenases herein. P450s are a superfamily of enzymes capable of catalyzing a wide variety of reactions including epoxidation, hydroxylation, O-dealkylations, desaturation etc. As discussed herein, one way of eliminating the toxicity of AF and ST, trichothecenes (T) and fumonisins (F) is to shuffle and select for a monooxygenase such as P450 which is capable of oxidation of mycotoxins. In one preferred embodiment, this monooxygenase nucleic acid is transduced into crop plants to make the plants mycotoxin resistant. With respect to AF, oxidation at the 15,16 position provides detoxification. Oxidative changes in other positions (for example, O-demethylation of the methoxy group in position 8, *see*, Silva et al. 1996) also help render the molecule nontoxic.

45 One particularly preferred source of p450 nucleic acids for shuffling is the *cyp* 1, 2 and 3 families of genes, e.g., from humans. *See*, <http://drnelson.utmem.edu/homepage.html>. A feature of the invention is the discovery

5 other applications, recombinant segments are extracted from the cells, and optionally
packaged as viruses, before screening.

10 The nature of screening or selection depends on what property or
characteristic is to be acquired or the property or characteristic for which improvement is
5 sought, and many examples are discussed below. It is not usually necessary to understand
the molecular basis by which particular products of recombination (recombinant
segments) have acquired new or improved properties or characteristics relative to the
15 starting substrates. For example, a mycotoxin detoxification gene can have many
component sequences each having a different intended role (e.g., coding sequence,
10 regulatory sequences, targeting sequences, stability-conferring sequences, subunit
sequences and sequences affecting integration). Each of these component sequences can
be varied and recombined simultaneously. Screening/selection can then be performed,
20 for example, for recombinant segments that have increased ability to confer mycotoxin
detoxification activity upon a cell without the need to attribute such improvement to any
25 of the individual component sequences of the vector.

Depending on the particular screening protocol used for a desired
property, initial round(s) of screening can sometimes be performed using bacterial cells
30 due to high transfection efficiencies and ease of culture. However, especially for
eukaryotic mycotoxin detoxification enzymes such as eukaryotic P450s, yeast, fungal or
20 other eukaryotic systems are optionally used for library expression and screening.
Similarly other types of screening which are not amenable to screening in bacterial or
35 simple eukaryotic library cells, are performed in cells selected for use in an environment
close to that of their intended use. Final rounds of screening can be performed in the
precise cell type of intended use.

40 25 If further improvement in a property is desired, at least one, and usually a
collection, of recombinant segments surviving a first round of screening/selection are
subject to a further round of recombination. These recombinant segments can be
recombined with each other or with exogenous segments representing the original
45 substrates or further variants thereof. Again, recombination can proceed *in vitro* or *in*
30 *vivo*. If the previous screening step identifies desired recombinant segments as
components of cells, the components can be subjected to further recombination *in vivo*,

5 Mullis *et al.*, (1987) U.S. Patent No. 4,683,202; *PCR Protocols A Guide to Methods and Applications* (Innis *et al.* eds) Academic Press Inc. San Diego, CA (1990) (Innis);
10 Arnheim & Levinson (October 1, 1990) *C&EN* 36-47; *The Journal Of NIH Research* (1991) 3, 81-94; (Kwoh *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86, 1173; Guatelli *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87, 1874; Lomell *et al.* (1989) *J. Clin. Chem* 35, 1826; Landegren *et al.*, (1988) *Science* 241, 1077-1080; Van Brunt (1990) *Biotechnology* 8, 291-294; Wu and Wallace, (1989) *Gene* 4, 560; Barringer *et al.* (1990) *Gene* 89, 117, and Sooknanan and Malek (1995) *Biotechnology* 13: 563-564. Improved methods of cloning *in vitro* amplified nucleic acids are described in Wallace *et al.*, U.S. Pat. No. 10 5,426,039. Improved methods of amplifying large nucleic acids by PCR are summarized in Cheng *et al.* (1994) *Nature* 369: 684-685 and the references therein, in which PCR amplicons of up to 40kb are generated. One of skill will appreciate that essentially any RNA can be converted into a double stranded DNA suitable for restriction digestion, PCR expansion and sequencing using reverse transcriptase and a polymerase. See, 15 Ausbel, Sambrook and Berger, *all supra*.

Oligonucleotides for use as probes, *e.g.*, in *in vitro* amplification methods, for use as gene probes, or as shuffling targets (*e.g.*, synthetic genes or gene segments) are typically synthesized chemically according to the solid phase phosphoramidite triester method described by Beaucage and Caruthers (1981), *Tetrahedron Letts.*, 20 22(20):1859-1862, *e.g.*, using an automated synthesizer, as described in Needham-VanDevanter *et al.* (1984) *Nucleic Acids Res.*, 12:6159-6168. 35 Oligonucleotides can also be custom made and ordered from a variety of commercial sources known to persons of skill.

Indeed, essentially any nucleic acid with a known sequence can be custom 40 ordered from any of a variety of commercial sources, such as The Midland Certified Reagent Company (mcrc@oligos.com), The Great American Gene Company (<http://www.genco.com>), ExpressGen Inc. (www.expressgen.com), Operon Technologies Inc. (Alameda, CA) and many others. Similarly, peptides and antibodies 45 can be custom ordered from any of a variety of sources, such as PeptidoGenic (pkim@ccnet.com), HTI Bio-products, inc. (<http://www.htibio.com>), BMA Biomedicals Ltd (U.K.), Bio Synthesis, Inc., and many others. 50

5 6254. Another example is Versicolorin B synthase, which synthesizes the side chain
cyclization of racemic versiconal hemiacetal to the bisfuran ring system of versicolorin B.
The dihydrobisfuran is important to the mutagenic nature of AFB₁ and ST. The isolation
10 and characterization of the Versicolorin B synthase gene from *Aspergillus parasiticus* and
5 the partial characterization of the related synthetic cluster is described in Silva (1996)
JBC 271(23):13600-13608. A total of twenty five co-regulated transcripts defining a ST
gene cluster, and containing most, or all, of the genes necessary for ST biosynthesis in *A.*
15 *nidulans* is described by Brown et al. (1996) PNAS 93:1418-1422. These genes are also
targets for shuffling for mycotoxin detoxification. Whole genome shuffling approaches
10 (described below) can also be used to select for plant cells which produce products that
20 down-regulate production of genes such as those described in Brown, *id.* (which provide
for ST and AF biosynthesis) thereby reducing ST and AF levels in target plants.

A variety of organisms known to contain additional monooxygenases
which could be shuffled in the methods of the invention are also known. The most
25 15 comprehensive studies on bacterial alkene epoxidation have been done on *Pseudomonas*
oleovorans. Work on *P. oleovorans* by May and coworkers (J. Biol. Chem., 248, 1725-
1730) showed that the monooxygenase contained in the cells is capable of epoxidizing
30 octene to 1,2-epoxy-octane in 70% enantiomeric purity. In addition, this enzyme is
capable of converting 1,7-octadiene to the diepoxide (May et al, J. Am. Chem. Soc., 98,
20 7856-7858) and 1,5-hexadiene and 1,11-dodecadiene to epoxides. This enzyme system is
also capable of mediating hydroxylation of longer chain alkanes (octanes, etc.) and fatty
35 acids. The enzyme has been cloned and sequenced and is comprised of three protein
components: rubredoxin (mw 19,000), NADH-rubredoxin reductase, and the ω -
hydroxylase (a non-heme iron protein). Microorganisms having MMO enzyme activities
40 25 with similar properties include the genera *Rhodococcus*, *Mycobacterium*, *Nocardia*
(*Nocardia carollina* B-276) and *Pseudomonas Corynebacterium equi* (IFO 3730). All of
these strains are available from ATCC and serve as sources for the genes which can be
isolated by hybridization and gene amplification methods.

45 30 Mycotoxin detoxification screening is done most easily in yeast, but a
bacterial system could also be constructed by co-expressing the accessory electron
transport proteins adrenodoxin and adrenodoxin reductase. DNA from clones with

improved activity can be shuffled together in subsequent rounds of DNA shuffling and screened for further improvement.

Assays for Mycotoxin Inactivation

Screening a number of cloned cytochrome P450 monooxygenases for activity vs. various AF and ST and other mycotoxins yields P450 nucleic acids specifying these reactions and other oxidative changes in 15, or 16 positions (like hydroxylation). These P450 genes can be of fungal, microbial, plant, insect or mammalian origin. The screen can be conducted by measuring the activity against any particular mycotoxin or against multiple mycotoxins, e.g., by preparing extracts of clones expressing P450 genes. The expected product(s) from the above toxins as well as other oxidized products derived from a P450 can be identified based on differences in physical properties (oxidation of mycotoxins causes a detectable difference in the physical characteristics of mycotoxins).

It is possible to directly select the clones expressing P450 specifying either specific or broad-based oxidation by using yeast, if the yeast are susceptible to the compound. For example, as noted above, Kimura et al (1997) describe expression of the *tri101* gene in yeast and selection of the yeast in medium containing T-2 toxin, a potent mycotoxin. This same assay format can be used for any mycotoxin which is toxic to yeast, or inhibitory to yeast growth on a medium.

Similarly, such assays can be performed using any of a variety of other cultured cells, by growing the cells (e.g., prokaryotic or eukaryotic cells) in the presence of a mycotoxin. To gradually select for more and more potent mycotoxin detoxification nucleic acids, cells are grown in medium containing increasingly high doses of the mycotoxin, e.g., following each round of a reiterative DNA shuffling procedure, as described herein.

In general, the culture of cells, including yeast, animal cells, plant cells and the like are well known. In addition to Berger, Ausubel and Sambrook, *all supra*, details on animal cell culture can be found in Freshney (*Culture of Animal Cells, a Manual of Basic Technique, third edition* Wiley-Liss, New York (1994)) and the references cited therein. The culture of plant cells is described, e.g., in Payne et al. (1992) Plant Cell and Tissue Culture in Liquid Systems John Wiley & Sons, Inc. New York, NY (Payne); and Gamborg and Phillips (eds) (1995) Plant Cell, Tissue and Organ

5 will be counted). In a variation of assay, with non-bead scintillating material, surfaces are activated with thiols or amines. All these variations are, in essence, SPA assays.

10 For Glutathione S-transferases (GSTs), the epoxide moiety of T2 is amenable to nucleophilic attack by thiol nucleophiles, including glutathione, whether transferred or not by GST. The thiol-T2 conjugate compound can be formed in an irreversible manner and is not an active toxin. Endogenous GST levels in plants are likely to be sufficient. Selection in yeast is used to evolve and screen as above for 3-OAT. DNA shuffling is used to optimize the specificity of plant GST enzymes towards T2 epoxide.

10 For epoxide hydrolase or isomerase it is sufficient to disrupt the T2 toxophore by modifying 12,13-epoxide to a glycol or an aldehyde. No known natural enzymes work on this epoxide. However, DNA shuffling is used to impart and optimize the required specificity. Selection in yeast is used to evolve and screen as for 3-OAT above.

15 Other assays for shuffling include chemical assays based on reactivity of residual epoxide, or formed rearranged 13-aldehyde product. One option is the use of a cytochrome P450 enzyme for aflatoxin detoxification by epoxidation of the double bond of the dihydrobisfuran moiety. Also, this can be used in conjunction with nucleophilic opening of the 15,16-epoxide (epoxide hydrolase, or GST, or an amine nucleophile, e.g. nucleobase or amino acid). Although 13-acetal can, in principle, be a subject to hydrolytic opening (enzymatic), the spontaneous toxophore regeneration may occur as it is favored by stereochemical means. Reactivity of 15,16-epoxide towards nucleophiles can be used for screen of P450s with the best activity towards AFB1. Exogenously supplied nucleophiles convenient for detection of AFB-epoxide-nucleophile adduct can be supplied in order to have shuffling done in bacterial or other microbial host which is insensitive to AFB1.

25 An alternative method for the assay of P450s with optimized activity towards aflatoxinB1 can use a variation of scintillation proximity assay using beads or other SPA material activated with a suitable nucleophilic group (amine, thiol) to trap any AFB1 15,16-epoxide. This uses a radioactively labeled AFB1 as a screen substrate. The latter can be prepared by chemical means, or by means of biosynthesis (with AFB1

5 producing *Aspergillus* strains) using a radioactively labeled AFB1 precursor/intermediate of its biosynthesis pathway.

10 Other microbial genes specifying oxygenases suitable for oxidation of aflatoxin such as AFB1 are cis-diol making dioxygenases with specificity for furan substructures. Other genes coding for flavoprotein monooxygenases of microbial or
5 mammalian origin are suitable candidates for shuffling for generating and improving activity vs. AF, ST and T. An way of degrading AFB1 is by opening the coumarin-lactone ring - C1 ester with a hydrolase. This is of value when the ring opening product is intercepted with a peroxidase for oxidative radical coupling of the free C12 phenol.
15 The toxophore is destroyed after this oxidation. Even without peroxidase-induced radical oxidation process, the hydrolytic option is of use because the lactone ring opening product is subject to spontaneous C-1 hydrolytic decarboxylation (beta-ketoacid).
20

As is apparent from the foregoing, the relevant assay will depend on the application. Many assays formats are suitable. Advantageously, any of the assays can be
25 practiced in a high-throughput format.
15

In the high throughput assays of the invention, it is possible to screen up to several thousand different shuffled variants in a single day. For example, each well of a
30 microtiter plate can be used to run a separate assay, or, if concentration or incubation time effects are to be observed, every 5-10 wells can test a single variant. Thus, a single standard microtiter plate can assay about 100 (e.g., 96) mycotoxin detoxification
20 reactions. If 1536 well plates are used, then a single plate can easily assay from about 100- about 1500 different reactions. It is possible to assay several different plates per day; assay screens for up to about 6,000-20,000 different assays (i.e., involving different
35 nucleic acids, encoded proteins, concentrations, etc.) is possible using the integrated systems of the invention. More recently, microfluidic approaches to reagent
40 manipulation have been developed, e.g., by Caliper Technologies (Palo Alto, CA).
25

In addition to fluidic approaches, it is possible simply to grow cells on plates of agar which comprise mycotoxins. Cells which have mycotoxin detoxification
45 activity (e.g., due to transduction with mycotoxin detoxification nucleic acids) are able to grow on the plates. This approach offers a very simple and high-throughput screening
30 method.

5

10

15

20

25

30

35

40

45

50

55

In one aspect, library members, e.g., cells, viral plaques, spores or the like, are separated on solid media to produce individual colonies (or plaques). Using an automated colony picker (e.g., the Q-bot, Genetix, U.K.), colonies or plaques are identified, picked, and up to 10,000 different mutants inoculated into 96 well microtitre dishes, optionally containing glass balls in the wells to prevent aggregation. The Q-bot does not pick an entire colony but rather inserts a pin through the center of the colony and exits with a small sampling of cells, (or mycelia) and spores (or viruses in plaque applications). The time the pin is in the colony, the number of dips to inoculate the culture medium, and the time the pin is in that medium each effect inoculum size, and each can be controlled and optimized. The uniform process of the Q-bot decreases human handling error and increases the rate of establishing cultures (roughly 10,000/4 hours). These cultures are then shaken in a temperature and humidity controlled incubator. The glass balls in the microtiter plates act to promote uniform aeration of cells dispersal of mycelial fragments, or the like, similar to the blades of a fermenter. Clones from cultures of interest can be cloned by limiting dilution. As also described supra, plaques or cells constituting libraries can also be screened directly for production of proteins, either by detecting hybridization, protein activity, protein binding to antibodies, or the like.

The ability to detect a subtle increase in the performance of a shuffled library member over that of a parent strain relies on the sensitivity of the assay. The chance of finding the organisms having an improvement in mycotoxin detoxification activity is increased by the number of individual mutants that can be screened by the assay. To increase the chances of identifying a pool of sufficient size, a prescreen that increases the number of mutants processed by 10-fold can be used. The goal of the primary screen will be to quickly identify mutants having equal or better product titres than the parent strain(s) and to move only these mutants forward to liquid cell culture for subsequent analysis.

A number of well known robotic systems have also been developed for solution phase chemistries useful in assay systems. These systems include automated workstations like the automated synthesis apparatus developed by Takeda Chemical Industries, LTD. (Osaka, Japan) and many robotic systems utilizing robotic arms

5 (Zymate II, Zymark Corporation, Hopkinton, Mass.; Orca, Hewlett-Packard, Palo Alto, Calif.) which mimic the manual synthetic operations performed by a scientist. Any of
10 the above devices are suitable for use with the present invention, e.g., for high-throughput screening of molecules encoded by codon-altered nucleic acids. The nature
5 and implementation of modifications to these devices (if any) so that they can operate as discussed herein with reference to the integrated system will be apparent to persons
15 skilled in the relevant art.

High throughput screening systems are commercially available (*see, e.g.,* Zymark Corp., Hopkinton, MA; Air Technical Industries, Mentor, OH; Beckman
10 Instruments, Inc. Fullerton, CA; Precision Systems, Inc., Natick, MA, *etc.*). These systems typically automate entire procedures including all sample and reagent pipetting,
20 liquid dispensing, timed incubations, and final readings of the microplate in detector(s) appropriate for the assay. These configurable systems provide high throughput and rapid
25 start up as well as a high degree of flexibility and customization.

15 The manufacturers of such systems provide detailed protocols the various high throughput. Thus, for example, Zymark Corp. provides technical bulletins
describing screening systems for detecting the modulation of gene transcription, ligand
30 binding, and the like. Microfluidic approaches to reagent manipulation have also been developed, e.g., by Caliper Technologies (Palo Alto, CA).

20 Optical images viewed (and, optionally, recorded) by a camera or other recording device (*e.g.,* a photodiode and data storage device) are optionally further
35 processed in any of the embodiments herein, *e.g.,* by digitizing the image and/or storing and analyzing the image on a computer. As noted above, in some applications,
40 mycotoxin detoxification products are florescent, making optical detection approaches
25 appropriate in these instances. A variety of commercially available peripheral equipment and software is available for digitizing, storing and analyzing a digitized video or
digitized optical image, *e.g.,* using PC (Intel x86 or pentium chip- compatible DOS™, OS2™ WINDOWS™, WINDOWS NT™ or WINDOWS95™ based machines),
45 MACINTOSH™, or UNIX based (*e.g.,* SUN™ work station) computers.

30 One conventional system carries light from the assay device to a cooled charge-coupled device (CCD) camera, in common use in the art. A CCD camera
50

5 includes an array of picture elements (pixels). The light from the specimen is imaged on
the CCD. Particular pixels corresponding to regions of the specimen (e.g., individual
10 hybridization sites on an array of biological polymers) are sampled to obtain light
intensity readings for each position. Multiple pixels are processed in parallel to increase
5 speed. The apparatus and methods of the invention are easily used for viewing any
sample, e.g., by fluorescent or dark field microscopic techniques.

15 Integrated systems for analysis in the present invention typically include a
digital computer with high-throughput liquid control software, image analysis software,
data interpretation software, a robotic liquid control armature for transferring solutions
10 from a source to a destination operably linked to the digital computer, an input device
(e.g., a computer keyboard) for entering data to the digital computer to control high
20 throughput liquid transfer by the robotic liquid control armature and, optionally, an
image scanner for digitizing label signals from labeled assay component. The image
scanner interfaces with the image analysis software to provide a measurement of optical
25 intensity. Typically, the intensity measurement is interpreted by the data interpretation
15 software to show whether the mycotoxin detoxification products are produced.

30 Monooxygenase activity can also be monitored by HPLC, gas
chromatography and mass spectroscopy, as well as a variety of other analytical methods
available to one of skill. Incorporation of ^{18}O from radio-labeled molecular oxygen can
20 be monitored directly by mass shift by MS methods and by an appropriate radioisotope
detector with HPLC and GC devices. In a high-throughput modality, a method of choice
35 is high-throughput MS, or MS with an electron spray-based detection method.

40 In addition, epoxide formation can be indirectly measured by various
reactive colorimetric reactions. For example, disappearance of peroxide over time can
25 be monitored directly either potentiometrically or colorimetrically using a number of
commercially available peroxide reactive dyes.

45 In one set of assays, the relative toxicity of mycotoxin products produced
by modification of mycotoxin detoxification enzymes is determined. In particular,
toxicity can be evaluated in any of the usual assays for mycotoxin toxicity and,
30 optionally, compared to the toxicity of the unmodified mycotoxin. In the event that
toxicity is reduced, secondary toxic effects of detoxification products can be evaluated
50

5 using these usual assays for mycotoxin activity, or using additional assays such as cell
survival assays, e.g., in the presence of increasing levels of the secondary product. This
secondary assay helps to determine which mycotoxin detoxification activities are most
10 desirable, i.e., using secondary toxicities of mycotoxin metabolites as a measure of
unwanted toxicity.

Formats for Nucleic Acid Recombination-Nucleic Acid Shuffling

15 The present invention involves shuffling of nucleic acids. The following
publications describe a variety of recursive recombination (shuffling) procedures and/or
methods which can be incorporated into such procedures: Stemmer, et al., (1999)
10 "Molecular breeding of viruses for targeting and other clinical properties. Tumor
Targeting" 4:1-4; Nasset al. (1999) "DNA Shuffling of subgenomic sequences of
20 subtilisin" Nature Biotechnology 17:893-896; Chang et al. (1999) "Evolution of a cytokine
using DNA family shuffling" Nature Biotechnology 17:793-797; Minshull and Stemmer
(1999) "Protein evolution by molecular breeding" Current Opinion in Chemical Biology
25 3:284-290; Christians et al. (1999) "Directed evolution of thymidine kinase for AZT
phosphorylation using DNA family shuffling" Nature Biotechnology 17:259-264;
Crameriet al. (1998) "DNA shuffling of a family of genes from diverse species accelerates
directed evolution" Nature 391:288-291; Crameri et al. (1997) "Molecular evolution of an
30 arsenate detoxification pathway by DNA shuffling," Nature Biotechnology 15:436-438;
Zhang et al. (1997) "Directed evolution of an effective fucosidase from a galactosidase by
DNA shuffling and screening" Proceedings of the National Academy of Sciences, U.S.A.
35 94:4504-4509; Patten et al. (1997) "Applications of DNA Shuffling to Pharmaceuticals and
Vaccines" Current Opinion in Biotechnology 8:724-733; Crameri et al. (1996)
"Construction and evolution of antibody-phage libraries by DNA shuffling" Nature
40 Medicine 2:100-103; Crameri et al. (1996) "Improved green fluorescent protein by
molecular evolution using DNA shuffling" Nature Biotechnology 14:315-319; Gates et al.
(1996) "Affinity selective isolation of ligands from peptide libraries through display on a
lac repressor 'headpiece dimer'" Journal of Molecular Biology 255:373-386; Stemmer
45 (1996) "Sexual PCR and Assembly PCR" In: The Encyclopedia of Molecular Biology.
VCH Publishers, New York. pp.447-457; Crameri and Stemmer (1995) "Combinatorial
multiple cassette mutagenesis creates all the permutations of mutant and wildtype
30 cassettes" BioTechniques 18:194-195; Stemmer et al., (1995) "Single-step assembly of a

5 AND REASSEMBLY" EP 0934999; Stemmer "EVOLVING CELLULAR DNA UPTAKE
BY RECURSIVE SEQUENCE RECOMBINATION" EP 0932670; Stemmer et al.,
10 "MODIFICATION OF VIRUS TROPISM AND HOST RANGE BY VIRAL GENOME
SHUFFLING" WO 9923107; Apt et al., "HUMAN PAPILLOMAVIRUS VECTORS" WO
5 9921979; Del Cardayre et al. "EVOLUTION OF WHOLE CELLS AND ORGANISMS
BY RECURSIVE SEQUENCE RECOMBINATION" WO 9831837; Patten and Stemmer,
15 "METHODS AND COMPOSITIONS FOR POLYPEPTIDE ENGINEERING" WO
9827230; Stemmer et al., and "METHODS FOR OPTIMIZATION OF GENE THERAPY
BY RECURSIVE SEQUENCE SHUFFLING AND SELECTION" WO9813487.

10 Certain U.S. Applications provide additional details regarding DNA
20 shuffling and related techniques, including "SHUFFLING OF CODON ALTERED
GENES" by Patten et al. filed September 29, 1998, (USSN 60/102,362), January 29, 1999
(USSN 60/117,729), and September 28, 1999, USSN _____ (Attorney Docket Number 20-
28520US/PCT); "EVOLUTION OF WHOLE CELLS AND ORGANISMS BY
25 RECURSIVE SEQUENCE RECOMBINATION", by del Cardyre et al. filed July 15, 1998
(USSN 09/166,188), and July 15, 1999 (USSN 09/354,922); "OLIGONUCLEOTIDE
MEDIATED NUCLEIC ACID RECOMBINATION" by Cramer et al., filed February 5,
30 1999 (USSN 60/118,813) and filed June 24, 1999 (USSN 60/141,049) and filed September
28, 1999 (USSN _____, Attorney Docket Number 02-29620US); and "USE OF CODON-
20 BASED OLIGONUCLEOTIDE SYNTHESIS FOR SYNTHETIC SHUFFLING" by
Welch et al., filed September 28, 1999 (USSN _____, Attorney Docket Number 02-
35 010070US); and "METHODS FOR MAKING CHARACTER STRINGS,
POLYNUCLEOTIDES & POLYPEPTIDES HAVING DESIRED CHARACTERISTICS"
by Selifonov and Stemmer, filed February 5, 1999 (USSN 60/118854).

40 25 As review of the foregoing publications, patents, published applications and
U.S. patent applications reveals, shuffling (or "recursive recombination") of nucleic acids
to provide new nucleic acids with desired properties can be carried out by a number of
45 methods. These methods can be adapted to the present invention to evolve the mycotoxin
detoxification activity as discussed herein to produce new mycotoxin detoxification nucleic
30 acids with new or improved properties. Both the methods of making such mycotoxin
detoxification nucleic acids and the mycotoxin detoxification nucleic acids produced by
these methods are a feature of the invention.

5

10

15

20

25

30

35

40

45

50

55

In brief, at least 5 different general classes of recombination methods are applicable to the present invention. First, nucleic acids can be recombined in vitro by any of a variety of techniques discussed in the references above, including e.g., DNase digestion of nucleic acids to be recombined followed by ligation and/or PCR reassembly of the nucleic acids. Second, nucleic acids can be recursively recombined in vivo, e.g., by allowing recombination to occur between nucleic acids in cells. Third, whole cell genome recombination methods can be used in which whole genomes of cells are recombined, optionally including spiking of the genomic recombination mixtures with desired library components such as mycotoxin detoxification nucleic acids homologue nucleic acids. Fourth, synthetic recombination methods can be used, in which oligonucleotides corresponding to different mycotoxin detoxification nucleic acid homologues are synthesized and reassembled in PCR or ligation reactions which include oligonucleotides which correspond to more than one parental nucleic acid, thereby generating new recombined nucleic acids. Oligonucleotides can be made by standard nucleotide addition methods, or can be made and shuffled by tri-nucleotide synthetic and shuffling approaches. Fifth, in silico methods of recombination can be effected in which genetic algorithms are used in a computer to recombine sequence strings which correspond to mycotoxin detoxification nucleic acid homologues. The resulting recombined sequence strings are optionally converted into nucleic acids by synthesis of nucleic acids which correspond to the recombined sequences, e.g., in concert with oligonucleotide synthesis/ gene reassembly techniques. Any of the preceding general recombination formats can be practiced in a reiterative fashion to generate a more diverse set of recombinant nucleic acids. In addition, these general approaches can, and often are, used in combination.

The above references provide these and other basic recombination formats as well as many modifications of these formats. Regardless of the format which is used, the nucleic acids of the invention can be recombined (with each other or with related (or even unrelated) nucleic acids to produce a diverse set of recombinant nucleic acids, including homologous nucleic acids. In general, the sequence recombination techniques described herein provide particular advantages in that they provide for recombination between mycotoxin detoxification nucleic acids, or derivatives thereof, in any available format, thereby providing a very fast way of exploring the manner in which different combinations of sequences can affect a desired result.

5 Following recombination, any nucleic acids which are produced can be selected for a desired activity. In the context of the present invention, this can include testing for and identifying any mycotoxin detoxification activities, by any of the assays in
10 the art. In addition, useful properties such as low crop yield enhancement, can also be simultaneously selected for. A variety of mycotoxin detoxification nucleic acid related (or even unrelated) properties can be assayed for, using any available assay.

15 A recombinant nucleic acid produced by recursively recombining one or more polynucleotide of the invention with one or more additional nucleic acid also forms a part of the invention. The one or more additional nucleic acid may include another
10 polynucleotide of the invention (i.e., one or more evolved mycotoxin detoxification nucleic acids); optionally, alternatively, or in addition, the one or more additional nucleic acid can include, e.g., a nucleic acid encoding a naturally-occurring mycotoxin detoxification
20 nucleic acid or a subsequence thereof, or any homologous sequence or subsequence thereof, or, e.g., any other homologous or non-homologous nucleic acid (certain
25 recombination formats noted above, notably those performed synthetically or in silico, do not require homology for recombination).

The recombining steps may be performed in vivo, in vitro, or in silico as described in more detail in the references above. Also included in the invention is a cell
30 containing any resulting recombinant nucleic acid, nucleic acid libraries produced by recursive recombination of the nucleic acids set forth herein, and populations of cells,
20 vectors, viruses, plasmids or the like comprising the library or comprising any recombinant nucleic acid resulting from recombination (or recursive recombination) of a nucleic acid as set forth herein with another such nucleic acid, or an additional nucleic acid.
35 Corresponding sequence strings in a database present in a computer system or computer readable medium are a feature of the invention.

40 Specific Formats for Sequence Recombination

DNA shuffling can be applied to a collection of mycotoxin without prior screening for activity vs. one or more mycotoxins. The shuffled genes can be cloned in
45 appropriate *E. coli* or yeast, and clones exhibiting desired activity can be selected as described above. The screening will be based e.g., on differences in the physical
30 properties between the parent mycotoxin and its modified, oxidized product, or upon cell survival on mycotoxin containing media. The final gene product can be optimized for

5 rapid oxidation and/or desired substrate specificity for one or more mycotoxins, by
further rounds of shuffling. The optimized gene or genes obtained after several rounds
10 of shuffling could be cloned into desired crops in which AF, ST or other mycotoxin
production by the appropriate pathogen is a problem. This will help eliminate the
5 toxicity associated with the pathogen-produced mycotoxins in the grains.

15 Shuffled gene(s) developed by the above screening method for the
identification of oxidation of one or more mycotoxins are optionally shuffled by at least
one of the five general approaches for sequence recombination noted above.

In specific formats, DNA-shuffling can be performed on a single gene.

20 Alternatively, several homologous genes can be identified by sequence comparison with
known homologous genes. These genes can be synthesized and shuffled as a family of
homologs, to select recombinants with the desired activity. The shuffled genes can be
cloned into *E. coli*, yeast, plants, fungi, or animal cells and those producing high activity
25 can be identified by the methods described above.

15 Whole genome shuffling can be performed to shuffle detoxification genes
(along with other genomic nucleic acids), thereby producing cells with enhanced
detoxification activity. For whole genome shuffling approaches, it is not even necessary
30 to identify which mycotoxin detoxification genes are being shuffled. Instead, e.g., plant
cell genomes are combined and shuffled to acquire mycotoxin detoxification activity, as
20 measured in any of the assays above.

35 Mycotoxin detoxification genes can be codon modified to access
mutational diversity not present in any naturally occurring detoxification gene. Shuffling
can be performed using synthetic shuffling and in silico approaches. Details on each of
40 these procedures can be found in the references noted above and as further detailed
25 below.

Generally, the methods of the invention entail performing DNA
recombination ("shuffling") and screening or selection to "evolve" individual genes,
45 whole plasmids or viruses, multigene clusters, or even whole genomes (e.g., Stemmer
(1995) *Bio/Technology* 13:549-553 and the other references noted herein). Reiterative
30 cycles of recombination and screening/selection can be performed to further evolve the
nucleic acids of interest. Such techniques do not require the extensive analysis and
50

5 computation required by conventional methods for polypeptide engineering. Shuffling
allows the recombination of large numbers of mutations in a minimum number of
10 selection cycles, in contrast to natural pair-wise recombination events (e.g., as occur
during sexual replication). Thus, the sequence recombination techniques described
5 herein provide particular advantages in that they provide recombination between
mutations in any or all of these, thereby providing a very fast way of exploring the
15 manner in which different combinations of mutations can affect a desired result. In some
instances, however, structural and/or functional information is available which, although
not required for sequence recombination, provides opportunities for modification of the
10 technique.

20 Exemplary formats and examples for sequence recombination, referred to,
e.g., as "DNA shuffling," "fast forced evolution," or "molecular breeding," have been
described by the present inventors and co-workers in the publications, patents and patent
25 applications noted above.

15 In one class of embodiments, the recombination procedure starts with at
least two substrates that generally show substantial sequence identity to each other (*i.e.*,
at least about 30%, 50%, 70%, 80% or 90% sequence identity), but differ from each
30 other at certain positions. The difference can be any type of mutation, for example,
substitutions, insertions and deletions. Often, different segments differ from each other
20 in about 5-20 positions. For recombination to generate increased diversity relative to the
starting materials, the starting materials must differ from each other in at least two
35 nucleotide positions. That is, if there are only two substrates, there should be at least
two divergent positions. If there are three substrates, for example, one substrate can
40 differ from the second at a single position, and the second can differ from the third at a
different single position. The starting DNA segments can be natural variants of each
25 other, for example, allelic or species variants. The segments can also be from nonallelic
genes showing some degree of structural and usually functional relatedness (*e.g.*,
45 different genes within a superfamily, such as the cytochrome P450 super family). The
starting DNA segments can also be induced variants of each other. For example, one
30 DNA segment can be produced by error-prone PCR replication of the other, or by
substitution of a mutagenic cassette. Induced mutants can also be prepared by
50

5 reassembly with DNA ligase. Typically, restriction enzyme (or endonuclease) sites are
identified by conventional restriction enzyme mapping techniques (Sambrook, Ausubel,
10 and Berger, *supra.*), by analysis of sequence information for that gene, or by
introduction of desired restriction sites into a nucleic acid sequence by synthesis (*i.e.* by
5 incorporation of silent mutations).

The DNA substrate molecules to be digested can either be from *in vivo*
15 replicated DNA, such as a plasmid preparation, or from PCR amplified nucleic acid
fragments harboring the restriction enzyme recognition sites of interest, preferably near
the ends of the fragment. Typically, at least two variants of a gene of interest, each
10 having one or more mutations, are digested with at least one restriction enzyme
determined to cut within the nucleic acid sequence of interest. The restriction fragments
are then joined with DNA ligase to generate full length genes having shuffled regions.
The number of regions shuffled will depend on the number of cuts within the nucleic acid
25 sequence of interest. The shuffled molecules can be introduced into cells as described
above and screened or selected for a desired property as described herein. Nucleic acid
15 can then be isolated from pools (libraries), or clones having desired properties and
subjected to the same procedure until a desired degree of improvement is obtained.

30 In some embodiments, at least one DNA substrate molecule or fragment
thereof is isolated and subjected to mutagenesis. In some embodiments, the pool or
20 library of religated restriction fragments are subjected to mutagenesis before the
digestion-ligation process is repeated. "Mutagenesis" as used herein comprises such
35 techniques known in the art as PCR mutagenesis, oligonucleotide-directed mutagenesis,
site-directed mutagenesis, etc., and recursive sequence recombination by any of the
techniques described herein.

40 25 *Reassembly PCR*

A further technique for recombining mutations in a nucleic acid sequence
utilizes "reassembly PCR." This method can be used to assemble multiple segments that
45 have been separately evolved into a full length nucleic acid template such as a gene. This
technique is performed when a pool of advantageous mutants is known from previous
30 work or has been identified by screening mutants that may have been created by any
mutagenesis technique known in the art, such as PCR mutagenesis, cassette mutagenesis,

5 doped oligo mutagenesis, chemical mutagenesis, or propagation of the DNA template *in*
10 *vivo* in mutator strains. Boundaries defining segments of a nucleic acid sequence of
interest preferably lie in intergenic regions, introns, or areas of a gene not likely to have
mutations of interest. Preferably, oligonucleotide primers (oligos) are synthesized for
5 PCR amplification of segments of the nucleic acid sequence of interest, such that the
sequences of the oligonucleotides overlap the junctions of two segments. The overlap
15 region is typically about 10 to 100 nucleotides in length. Each of the segments is
amplified with a set of such primers. The PCR products are then "reassembled"
according to assembly protocols such as those discussed herein to assemble randomly
10 fragmented genes. In brief, in an assembly protocol the PCR products are first purified
20 away from the primers, by, for example, gel electrophoresis or size exclusion
chromatography. Purified products are mixed together and subjected to about 1-10 cycles
of denaturing, reannealing, and extension in the presence of polymerase and
25 deoxynucleoside triphosphates (dNTP's) and appropriate buffer salts in the absence of
15 additional primers ("self-priming"). Subsequent PCR with primers flanking the gene are
used to amplify the yield of the fully reassembled and shuffled genes.

30 In some embodiments, the resulting reassembled genes are subjected to
mutagenesis before the process is repeated.

35 In a further embodiment, the PCR primers for amplification of segments
20 of the nucleic acid sequence of interest are used to introduce variation into the gene of
interest as follows. Mutations at sites of interest in a nucleic acid sequence are identified
by screening or selection, by sequencing homologues of the nucleic acid sequence, and so
on. Oligonucleotide PCR primers are then synthesized which encode wild type or mutant
40 information at sites of interest. These primers are then used in PCR mutagenesis to
25 generate libraries of full length genes encoding permutations of wild type and mutant
information at the designated positions. This technique is typically advantageous in cases
where the screening or selection process is expensive, cumbersome, or impractical
45 relative to the cost of sequencing the genes of mutants of interest and synthesizing
mutagenic oligonucleotides.

5

Oligonucleotide and in silico shuffling formats for Mycotoxin
Detoxification Shuffling

10

15

20

25

30

35

40

45

50

55

Two additional related formats are useful in the practice of the present invention. The first, referred to as "in silico" shuffling utilizes computer algorithms to perform "virtual" shuffling using genetic operators in a computer. As applied to the present invention, mycotoxin detoxification nucleic acid sequence strings are recombined in a computer system and desirable products are made, e.g., by reassembly PCR or ligation of synthetic oligonucleotides, or other available techniques. In silico shuffling is described in detail in Selifonov and Stemmer in "METHODS FOR MAKING CHARACTER STRINGS, POLYNUCLEOTIDES & POLYPEPTIDES HAVING DESIRED CHARACTERISTICS" filed 02/05/1999, USSN 60/118854. In brief, genetic operators (algorithms which represent given genetic events such as point mutations, recombination of two strands of homologous nucleic acids, etc.) are used to model recombinational or mutational events which can occur in one or more nucleic acid, e.g., by aligning nucleic acid sequence strings (using standard alignment software, or by manual inspection and alignment) and predicting recombinational outcomes based upon selected genetic algorithms (mutation, recombination, etc.). The predicted recombinational outcomes are used to produce corresponding molecules, e.g., by oligonucleotide synthesis and reassembly PCR. As applied to the present invention, mycotoxin detoxification nucleic acids are aligned and recombined in silico, using any desired genetic operator, to produce mycotoxin detoxification character strings which are then generated synthetically for subsequent screening.

The second useful format is referred to as "oligonucleotide mediated shuffling" in which oligonucleotides corresponding to a family of related homologous nucleic acids (e.g., as applied to the present invention, families of homologous mycotoxin detoxification variants of a nucleic acid) which are recombined to produce selectable nucleic acids. This format is described in detail in Crameri et al. "OLIGONUCLEOTIDE MEDIATED NUCLEIC ACID RECOMBINATION" filed February 5, 1999, USSN 60/118,813 and Crameri et al. "OLIGONUCLEOTIDE MEDIATED NUCLEIC ACID RECOMBINATION" filed June 24, 1999, USSN 60/141,049. In brief, selected oligonucleotides corresponding to multiple homologous parental nucleic acids are

5 synthesized, ligated and elongated (typically in a recursive format), typically either in a
polymerase or ligase-mediated elongation reaction, to produce full-length mycotoxin
detoxification nucleic acids. The technique can be used to recombine homologous or even
10 non-homologous mycotoxin detoxification nucleic acid sequences.

5 One advantage of oligonucleotide-mediated recombination is the ability to
recombine homologous nucleic acids with low sequence similarity, or even non-
homologous nucleic acids. In these low-homology oligonucleotide shuffling methods, one
15 or more set of fragmented nucleic acids (e.g., oligonucleotides corresponding to multiple
mycotoxin detoxification nucleic acids) are recombined, e.g., with a set of crossover
family diversity oligonucleotides. Each of these crossover oligonucleotides have a
20 plurality of sequence diversity domains corresponding to a plurality of sequence diversity
domains from homologous or non-homologous nucleic acids with low sequence similarity.
The fragmented oligonucleotides, which are derived by comparison to one or more
homologous or non-homologous nucleic acids, can hybridize to one or more region of the
25 crossover oligos, facilitating recombination.

15 When recombining homologous nucleic acids, sets of overlapping family
gene shuffling oligonucleotides (which are derived by comparison of homologous nucleic
acids, by synthesis of corresponding oligonucleotides) are hybridized and elongated (e.g.,
30 by reassembly PCR or ligation), providing a population of recombined nucleic acids,
which can be selected for a desired trait or property. The set of overlapping family
20 shuffling gene oligonucleotides includes a plurality of oligonucleotide member types
which have consensus region subsequences derived from a plurality of homologous target
35 nucleic acids.

Typically, as applied to the present invention, family gene shuffling
25 oligonucleotide which include one or more mycotoxin detoxification nucleic acid(s) are
provided by aligning homologous nucleic acid sequences to select conserved regions of
sequence identity and regions of sequence diversity. A plurality of family gene shuffling
oligonucleotides are synthesized (serially or in parallel) which correspond to at least one
45 region of sequence diversity.

30 Sets of fragments, or subsets of fragments used in oligonucleotide shuffling
approaches can be provided by cleaving one or more homologous nucleic acids (e.g., with
a DNase), or, more commonly, by synthesizing a set of oligonucleotides corresponding to a
50

5 plurality of regions of at least one nucleic acid (typically oligonucleotides corresponding to
a full-length nucleic acid are provided as members of a set of nucleic acid fragments). In
the shuffling procedures herein, these cleavage fragments can be used in conjunction with
10 family gene shuffling oligonucleotides, e.g., in one or more recombination reaction to
5 produce recombinant mycotoxin detoxification nucleic acid(s).

One final synthetic variant worth noting is found in "SHUFFLING OF
15 CODON ALTERED GENES" by Patten et al. filed September 29, 1998, (USSN
60/102,362), January 29, 1999 (USSN 60/117,729), and September 28, 1999,
USSN _____ (Attorney Docket Number 20-28520US/PCT). As noted in detail in this set
10 of related applications, one way of generating diversity in a set of nucleic acids to be
shuffled (i.e., as applied to the present invention, mycotoxin detoxification nucleic acids),
20 is to provide codon-altered nucleic acids which can be shuffled to provide access to
sequence space not present in naturally occurring sequences. In brief, by synthesizing
nucleic acids in which the codons which encode polypeptides are altered, it is possible to
25 access a completely different mutational spectrum upon subsequent mutation of the
nucleic acid. This increases the sequence diversity of the starting nucleic acids for
shuffling protocols, which alters the rate and results of forced evolution procedures.
30 Codon modification procedures can be used to modify any mycotoxin detoxification
nucleic acid herein, e.g., prior to performing DNA shuffling.

20 In brief, oligonucleotide sets comprising codon variations are synthesized
and reassembled into full-length nucleic acids. The oligonucleotide sets can themselves be
35 shuffled (e.g., where the oligonucleotides to be reassembled provide sequence diversity at
selected sites), and/or the full-length sequences can be shuffled by any available procedure
to produce diverse sets of mycotoxin detoxification nucleic acids.

40 25 *Site Directed Mutagenesis (SDM) with Oligonucleotides Encoding Homologue
Mutations Followed by Shuffling*

In some embodiments of the invention, sequence information from one or
more substrate sequences is added to a given "parental" sequence of interest, with
45 subsequent recombination between rounds of screening or selection. Typically, this is
30 done with site-directed mutagenesis performed by techniques well known in the art (e.g.,
Berger, Ausubel and Sambrook, *supra.*), or by the oligonucleotide or in silico methods

5 noted above, with one substrate as template and oligonucleotides encoding single or
multiple mutations from other substrate sequences, *e.g.* homologous genes. After
10 screening or selection for an improved phenotype of interest, the selected recombinant(s)
can be further evolved using RSR techniques described herein. After screening or
5 selection, site-directed mutagenesis can be done again with another collection of
oligonucleotides encoding homologous mutations, and the above process repeated until the
15 desired properties are obtained.

When the difference between two homologues is one or more single point
mutations in a codon, degenerate oligonucleotides can be used that encode the sequences
10 in both homologues. One oligonucleotide can include many such degenerate codons and
still allow one to exhaustively search all permutations over that block of sequence.

When the homologue sequence space is very large, it can be advantageous
to restrict the search to certain variants. Thus, for example, computer modeling tools
25 (Lathrop *et al.* (1996) *J. Mol. Biol.*, 255: 641-665) can be used to model each homologue
15 mutation onto the target protein and discard any mutations that are predicted to grossly
disrupt structure and function.

30 *In Vitro DNA Shuffling Formats*

In one embodiment for shuffling DNA sequences *in vitro*, the initial
substrates for recombination are a pool of related sequences, *e.g.*, different variant
20 forms, as homologs from different individuals, strains, or species of an organism, or
related sequences from the same organism, as allelic variations. The sequences can be
35 DNA or RNA and can be of various lengths depending on the size of the gene or DNA
fragment to be recombined or reassembled. Preferably the sequences are from 40 base
pairs (bp) to 50 kilobases (kb).

25 The pool of related substrates are converted into overlapping fragments,
e.g., from about 5 bp to 5 kb or more. Often, for example, the size of the fragments is
from about 10 bp to 1000 bp, *e.g.*, about 30 or 40 bp to about 100bp, *e.g.*, about 100 bp
45 to 500 bp. The conversion can be effected by a number of different methods, such as
DNase I or RNase digestion, random shearing, partial restriction enzyme digestion or
30 oligonucleotide synthesis. For discussions of protocols for the isolation, manipulation,
enzymatic digestion, and the like of nucleic acids, see, for example, Sambrook *et al.* and
50

Ausubel, both *supra*. The concentration of nucleic acid fragments of a particular length and sequence is often less than 0.1 % or 1 % by weight of the total nucleic acid. The number of different specific nucleic acid fragments in the mixture is usually at least about 100, 500 or 1000.

The mixed population of nucleic acid fragments are converted to at least partially single-stranded form using a variety of techniques, including, for example, heating, chemical denaturation, use of DNA binding proteins, and the like. Conversion can be effected by heating to about 80°C to 100°C, more preferably from 90°C to 96°C, to form single-stranded nucleic acid fragments and then reannealing. Conversion can also be effected by treatment with single-stranded DNA binding protein (see Wold (1997) *Annu. Rev. Biochem.* 66:61-92) or *recA* protein (see, e.g., Kitanitsa (1997) *Proc. Natl. Acad. Sci. U S A* 94:7837-7840). Single-stranded nucleic acid fragments having regions of sequence identity with other single-stranded nucleic acid fragments can then be reannealed by cooling to 20°C to 75°C, and preferably from 40°C to 65°C.

Renaturation can be accelerated by the addition of polyethylene glycol (PEG), other volume-excluding reagents or salt. The salt concentration is preferably from 0 mM to 200 mM, more preferably the salt concentration is from 10 mM to 100 mM. The salt may be KCl or NaCl. The concentration of PEG is preferably from 0% to 20%, more preferably from 5% to 10%. The fragments that reanneal can be from different substrates. The annealed nucleic acid fragments are incubated in the presence of a nucleic acid polymerase, such as Taq or Klenow, and dNTP's (*i.e.* dATP, dCTP, dGTP and dTTP). If regions of sequence identity are large, Taq polymerase can be used with an annealing temperature of between 45-65°C. If the areas of identity are small, Klenow polymerase can be used with an annealing temperature of between 20-30°C. The polymerase can be added to the random nucleic acid fragments prior to annealing, simultaneously with annealing or after annealing.

The process of denaturation, renaturation and incubation in the presence of polymerase of overlapping fragments to generate a collection of polynucleotides containing different permutations of fragments is sometimes referred to as shuffling of the nucleic acid *in vitro*. This cycle is optionally repeated for a desired number of times.

5

10

15

Preferably the cycle is repeated from 2 to 100 times, more preferably the sequence is repeated from 10 to 40 times. The resulting nucleic acids are a selectable family of double-stranded polynucleotides of from about 50 bp to about 100 kb, preferably from 500 bp to 50 kb. The population represents variants of the starting substrates showing substantial sequence identity thereto but also diverging at several positions. The population has many more members than the starting substrates. The population of fragments resulting from shuffling is used to transform host cells, optionally after cloning into a vector.

20

25

30

35

In one embodiment utilizing *in vitro* shuffling, subsequences of recombination substrates can be generated by amplifying the full-length sequences under conditions which produce a substantial fraction, typically at least 20 percent or more, of incompletely extended amplification products. Another embodiment uses random primers to prime the entire template DNA to generate less than full length amplification products. The amplification products, including the incompletely extended amplification products are denatured and subjected to at least one additional cycle of reannealing and amplification. This variation, in which at least one cycle of reannealing and amplification provides a substantial fraction of incompletely extended products, is termed "stuttering." In the subsequent amplification round, the partially extended (less than full length) products reanneal to and prime extension on different sequence-related template species. In another embodiment, the conversion of substrates to fragments can be effected by partial PCR amplification of substrates.

40

45

50

In another embodiment, a mixture of fragments is spiked with one or more oligonucleotides. The oligonucleotides can be designed to include precharacterized mutations of a wildtype sequence, or sites of natural variations between individuals or species. The oligonucleotides also include sufficient sequence or structural homology flanking such mutations or variations to allow annealing with the wildtype fragments. Annealing temperatures can be adjusted depending on the length of homology.

In a further embodiment, recombination occurs in at least one cycle by template switching, such as when a DNA fragment derived from one template primes on the homologous position of a related but different template. Template switching can be induced by addition of recA (*see*, Kiiianitsa (1997) *supra*), rad51 (*see*, Namsaraev (1997)

55

5 *Mol. Cell. Biol.* 17:5359-5368), *rad55* (see, Clever (1997) *EMBO J.* 16:2535-2544),
10 *rad57* (see, Sung (1997) *Genes Dev.* 11:1111-1121) or other polymerases (e.g., viral
polymerases, reverse transcriptase) to the amplification mixture. Template switching can
also be increased by increasing the DNA template concentration.

5 Another embodiment utilizes at least one cycle of amplification, which can
be conducted using a collection of overlapping single-stranded DNA fragments of related
15 sequence, and different lengths. Fragments can be prepared using a single stranded DNA
phage, such as M13 (see, Wang (1997) *Biochemistry* 36:9486-9492). Each fragment can
hybridize to and prime polynucleotide chain extension of a second fragment from the
10 collection, thus forming sequence-recombined polynucleotides. In a further variation,
ssDNA fragments of variable length can be generated from a single primer by Pfu, Taq,
20 Vent, Deep Vent, UITma DNA polymerase or other DNA polymerases on a first DNA
template (see, Cline (1996) *Nucleic Acids Res.* 24:3546-3551). The single stranded DNA
fragments are used as primers for a second, Kunkel-type template, consisting of a
25 uracil-containing circular ssDNA. This results in multiple substitutions of the first
template into the second. See, Levichkin (1995) *Mol. Biology* 29:572-577; Jung (1992)
30 *Gene* 121:17-24.

30 In some embodiments of the invention, shuffled nucleic acids obtained by
use of the recursive recombination methods of the invention, are put into a cell and/or
20 organism for screening. Shuffled monooxygenase genes can be introduced into, for
example, bacterial cells, yeast cells, fungal cells vertebrate cells, invertebrate cells or
35 plant cells for initial screening. *Bacillus* species (such as *B. subtilis* and *E. coli* are two
examples of suitable bacterial cells into which one can insert and express shuffled
monooxygenase genes which provide for convenient shuttling to other cell types (a
40 variety of vectors for shuttling material between these bacterial cells and eukaryotic cells
are available; see, Sambrook, Ausubel and Berger, *all supra*). The shuffled genes can be
25 introduced into bacterial, fungal or yeast cells either by integration into the chromosomal
DNA or as plasmids.

45 Although bacterial and yeast systems are most preferred in the present
30 invention, in one embodiment, shuffled genes can also be introduced into plant cells for
production purposes (it will be appreciated that transgenic plants are, increasingly, an
50

5 of multiple genome copies in such cells results in a higher frequency of homologous
recombination in these cells, both between copies of a gene in different genomes within
10 the cell, and between a genome within the cell and a transfected fragment. The increased
frequency of recombination allows one to evolve a gene evolved more quickly to acquire
5 optimized characteristics.

15 In nature, the existence of multiple genomic copies in a cell type would
usually not be advantageous due to the greater nutritional requirements needed to
maintain this copy number. However, artificial conditions can be devised to select for
high copy number. Modified cells having recombinant genomes are grown in rich media
20 (in which conditions, multicopy number should not be a disadvantage) and exposed to a
mutagen, such as ultraviolet or gamma irradiation or a chemical mutagen, *e.g.*,
mitomycin, nitrous acid, photoactivated psoralens, alone or in combination, which
induces DNA breaks amenable to repair by recombination. These conditions select for
25 cells having multicopy number due to the greater efficiency with which mutations can be
excised. Modified cells surviving exposure to mutagen are enriched for cells with
multiple genome copies. If desired, selected cells can be individually analyzed for
genome copy number (*e.g.*, by quantitative hybridization with appropriate controls). For
30 example, individual cells can be sorted using a cell sorter for those cells containing more
DNA, *e.g.*, using DNA specific fluorescent compounds or sorting for increased size
using light dispersion. Some or all of the collection of cells surviving selection are tested
20 for the presence of a gene that is optimized for the desired property.

35 In one embodiment, phage libraries are made and recombined in mutator
strains such as cells with mutant or impaired gene products of *mutS*, *mutT*, *mutH*, *mutL*,
ovrD, *dcm*, *vsr*, *umuC*, *umuD*, *sbcB*, *recJ*, etc. The impairment is achieved by genetic
40 mutation, allelic replacement, selective inhibition by an added reagent such as a small
25 compound or an expressed antisense RNA, or other techniques. High multiplicity of
infection (MOI) libraries are used to infect the cells to increase recombination frequency.

45 Additional strategies for making phage libraries and or for recombining
DNA from donor and recipient cells are set forth in U.S. Pat. No. 5,521,077.

30 Additional recombination strategies for recombining plasmids in yeast are set forth in
WO 97 07205.

5

Recursive macroshuffling techniques are described in U.S. Pat. 5,811,238 to Stemmer.

10

Whole Genome Shuffling

15

In one embodiment, the selection methods herein are utilized in a "whole genome shuffling" format. An extensive guide to the many forms of whole genome shuffling is found in the pioneering application to the inventors and their co-workers entitled "Evolution of Whole Cells and Organisms by Recursive Sequence Recombination," by del Cardyre et al. e.g., WO98/31837, PCT/US99/15972, and 09/354,922.

20

10 In brief, whole genome shuffling makes no presuppositions at all regarding what nucleic acids may confer a desired property. Instead, entire genomes (e.g., from a genomic library, or isolated from an organism) are shuffled in cells and selection protocols applied to the cells. Thus, as applied to the present invention, cell genomes or sub genomes (e.g., libraries) are recombined and resulting recombinant cells comprising the recombined nucleic acids are selected for mycotoxin detoxification activity. Thus, one feature of the invention is a cell (e.g., plant, animal, bacterial, or even fungal cell which comprises mycotoxin detoxification activity). These cells can be used to produce anti-mycotoxin extracts or can be applied to reduce fungal growth and/or pathogenicity in a system of interest.

25

30

20

Use of RecA

35

40

45

25

The frequency of recombination between nucleic acids in the mycotoxin detoxification shuffling procedures herein can be increased by coating the nucleic acids with a recombinogenic protein, e.g., before or after introduction into cells. See Pati et al., *Molecular Biology of Cancer* 1, 1 (1996); Sena & Zarling, *Nature Genetics* 3, 365 (1996); Revet et al., *J. Mol. Biol.* 232, 779-791 (1993); Kowalczykowski & Zarling in *Gene Targeting* (CRC 1995), Ch. 7. The recombinogenic protein promotes homologous pairing and/or strand exchange. The best characterized *recA* protein is from *E. coli* and is available from Pharmacia (Piscataway, NJ). In addition to the wild-type protein, a number of mutant *recA*-like proteins have been identified (e.g., *recA803*). Further, many organisms have *recA*-like recombinases with strand-transfer activities (e.g., Ogawa et al., *Cold Spring Harbor Symposium on Quantitative Biology* 18, 567-576 (1993);

50

55

Johnson & Symington, *Mol. Cell. Biol.* 15, 4843-4850 (1995); Fugisawa et al., *Nucl. Acids Res.* 13, 7473 (1985); Hsieh et al., *Cell* 44, 885 (1986); Hsieh et al., *J. Biol. Chem.* 264, 5089 (1989); Fishel et al., *Proc. Natl. Acad. Sci. USA* 85, 3683 (1988); Cassuto et al., *Mol. Gen. Genet.* 208, 10 (1987); Ganea et al., *Mol. Cell Biol.* 7, 3124 (1987); Moore et al., *J. Biol. Chem.* 19, 11108 (1990); Keene et al., *Nucl. Acids Res.* 12, 3057 (1984); Kimiec, *Cold Spring Harbor Symp.* 48, 675 (1984); Kimeic, *Cell* 44, 545 (1986); Kolodner et al., *Proc. Natl. Acad. Sci. USA* 84, 5560 (1987); Sugino et al., *Proc. Natl. Acad. Sci. USA* 85, 3683 (1985); Halbrook et al., *J. Biol. Chem.* 264, 21403 (1989); Eisen et al., *Proc. Natl. Acad. Sci. USA* 85, 7481 (1988); McCarthy et al., *Proc. Natl. Acad. Sci. USA* 85, 5854 (1988); Lowenhaupt et al., *J. Biol. Chem.* 264, 20568 (1989). Examples of such recombinase proteins include *recA*, *recA803*, *uvsX*, (Roca, A.I., *Crit. Rev. Biochem. Molec. Biol.* 25, 415 (1990)), *sepl* (Kolodner et al., *Proc. Natl. Acad. Sci. (U.S.A.)* 84, 5560 (1987); Tishkoff et al., *Molec. Cell. Biol.* 11, 2593), *RuvC* (Dunderdale et al., *Nature* 354, 506 (1991)), *DST2*, *KEM1*, *XRN1* (Dykstra et al., *Molec. Cell. Biol.* 11, 2583 (1991)), *STPα/DST1* (Clark et al., *Molec. Cell. Biol.* 11, 2576 (1991)), *HPP-1* (Moore et al., *Proc. Natl. Acad. Sci. (U.S.A.)* 88, 9067 (1991)), other eukaryotic recombinases (Bishop et al., *Cell* 69, 439 (1992); Shinohara et al., *Cell* 69, 457. *RecA* protein forms a nucleoprotein filament when it coats a single-stranded DNA. In this nucleoprotein filament, one monomer of *recA* protein is bound to about 3 nucleotides. This property of *recA* to coat single-stranded DNA is essentially sequence independent, although particular sequences favor initial loading of *recA* onto a polynucleotide (e.g., nucleation sequences). The nucleoprotein filament(s) can be formed on essentially any DNA to be shuffled and can form complexes with both single-stranded and double-stranded DNA in procaryotic and eukaryotic cells. *RecA* mediated techniques are also found in WO/93/22443.

Before contacting with *recA* or other recombinase, mycotoxin detoxification fragments are optionally denatured, e.g., by heat-treatment. *RecA* protein is then added at a concentration of about 1-10 μ M. After incubation, the *recA*-coated single-stranded DNA is introduced into recipient cells by conventional methods, such as chemical transformation or electroporation. In whole cell shuffling techniques, the fragments undergo homologous recombination with cognate endogenous genes. Because

of the increased frequency of recombination due to recombinase coating, the fragments need not be introduced as components of vectors.

Fragments are sometimes coated with other nucleic acid binding proteins that promote recombination, protect nucleic acids from degradation, or target nucleic acids to the nucleus. Examples of such proteins includes *Agrobacterium virE2* (Durrenberger et al., *Proc. Natl. Acad. Sci. USA* 86, 9154-9158 (1989)). Alternatively, recipient strains can be deficient in RecD activity. Single stranded ends can also be generated by 3'-5' exonuclease activity or restriction enzymes producing 5' overhangs.

Transducing Shuffled Nucleic Acids into Plants.

As noted herein, it is particularly desirable to transduce plants with shuffled nucleic acids to reduce the level of mycotoxins in the plants, and/or to practice the shuffling procedures in plant cells. Reduction of mycotoxins benefits the plants by making them resistant to mycotoxicosis, as well as be making the plants safer for consumption.

Methods of transducing plant cells with nucleic acids are generally available. In addition to Berger, Ausubel and Sambrook, useful general references for plant cell cloning, culture and regeneration include Jones (ed) (1995) Plant Gene Transfer and Expression Protocols-- Methods in Molecular Biology, Volume 49 Human Press Towata NJ; Payne et al. (1992) Plant Cell and Tissue Culture in Liquid Systems John Wiley & Sons, Inc. New York, NY (Payne); and Gamborg and Phillips (eds) (1995) Plant Cell, Tissue and Organ Culture: Fundamental Methods Springer Lab Manual, Springer-Verlag (Berlin Heidelberg New York) (Gamborg). A variety of Cell culture media are described in Atlas and Parks (eds) The Handbook of Microbiological Media (1993) CRC Press, Boca Raton, FL (Atlas). Additional information for plant cell culture is found in available commercial literature such as the Life Science Research Cell Culture Catalogue (1998) from Sigma- Aldrich, Inc (St Louis, MO) (Sigma-LSRCCC) and, e.g., the Plant Culture Catalogue and supplement (1997) also from Sigma-Aldrich, Inc (St Louis, MO) (Sigma-PCCS). Additional details regarding plant cell culture are found in R.R.D.Croy, Ed. Plant Molecular Biology (1993) Bios Scientific Publishers, Oxford, U.K.

The nucleic acid constructs of the invention are introduced into plant cells,

5 either in culture or in the organs of a plant by a variety of conventional techniques. For
example, the DNA construct can be introduced directly into the genomic DNA of the
10 plant cell using techniques such as electroporation and microinjection of plant cell
protoplasts, or the DNA constructs can be introduced directly to plant cells using ballistic
5 methods, such as DNA particle bombardment. Alternatively, the DNA constructs are
combined with suitable T-DNA flanking regions and introduced into a conventional
15 *Agrobacterium tumefaciens* host vector. The virulence functions of the *Agrobacterium*
tumefaciens host directs the insertion of the construct and adjacent marker into the plant
cell DNA when the cell is infected by the bacteria.

10 Microinjection techniques are known in the art and well described in the
scientific and patent literature. For example, a number of methods are described in Jones
20 (ed) (1995) Plant Gene Transfer and Expression Protocols-- Methods in Molecular
Biology, Volume 49 Human Press Towata NJ, as well as in the other references noted
herein and available in the literature.

15 For example, the introduction of DNA constructs using polyethylene
glycol precipitation is described in Paszkowski, *et al.*, EMBO J. 3:2717 (1984).
Electroporation techniques are described in Fromm, *et al.*, Proc. Nat'l. Acad. Sci. USA
30 82:5824 (1985). Ballistic transformation techniques are described in Klein, *et al.*, Nature
327:70-73 (1987). Additional details are found in Jones (1995) *supra*.

20 *Agrobacterium tumefaciens*-mediated transformation techniques, including
disarming and use of binary vectors, are also well described in the scientific literature.
35 See, for example Horsch, *et al.*, Science 233:496-498 (1984), and Fraley, *et al.*, Proc.
Nat'l. Acad. Sci. USA 80:4803 (1983). *Agrobacterium*-mediated transformation is a
preferred method of transformation of both monocots and particularly dicots.

40 25 To use shuffled sequences, recombinant DNA vectors suitable for
transformation of plant cells are prepared. A DNA sequence coding for the desired
shuffled mycotoxin detoxification DNA is transduced into the plant. Where the sequence
45 is expressed, the sequence is optionally combined with transcriptional and translational
initiation regulatory sequences which further direct the transcription or translation of the
30 sequence from shuffled the gene in the intended tissues of the transformed plant.

Regeneration of Transgenic Plants

Transformed plant cells which are derived by any of the above transformation techniques can be cultured to regenerate a whole plant which possesses the transformed genotype and thus the desired phenotype. Such regeneration techniques rely on manipulation of certain phytohormones in a tissue culture growth medium, typically relying on a biocide and/or herbicide marker which has been introduced together with the desired nucleotide sequences. Plant regeneration from cultured protoplasts is described in Evans, et al., Protoplasts Isolation and Culture, Handbook of Plant Cell Culture, pp. 124-176, Macmillian Publishing Company, New York, (1983); and Binding, Regeneration of Plants, Plant Protoplasts, pp. 21-73, CRC Press, Boca Raton, (1985). Regeneration can also be obtained from plant callus, explants, somatic embryos (Dandekar, et al., J. Tissue Cult. Meth. 12:145 (1989); McGranahan, et al., Plant Cell Rep. 8:512 (1990)), organs, or parts thereof. Such regeneration techniques are described generally in Klee, et al., Ann. Rev. of Plant Phys. 38:467-486 (1987). Additional details are found in Payne (1992) and Jones (1995), *both supra*.

Preferred plants for expression of mycotoxin resistance genes include those for which mycotoxins are a significant problem, such as plants in the family *Graminae* (including corn, rye, triticale, barley, millet, rice, wheat, oats, etc.) plants in the family *Leguminosa* (e.g., peanuts, peas, beans, and the like) and nut plants (walnut, pecan, etc.) .

More generally, important commercial crops which benefit from mycotoxin detoxification include both monocots and dicots. Monocots such as plants in the grass family (*Gramineae*), such as plants in the sub families *Fetuoideae* and *Poacoideae*, which together include several hundred genera including plants in the genera *Agrostis*, *Phleum*, *Dactylis*, *Sorgum*, *Setaria*, *Zea* (e.g., corn), *Oryza* (e.g., rice), *Triticum* (e.g., wheat), *Secale* (e.g., rye), *Avena* (e.g., oats), *Hordeum* (e.g., barley), *Saccharum*, *Poa*, *Festuca*, *Stenotaphrum*, *Cynodon*, *Coix*, the *Olyreae*, *Phareae* and many others. As noted, plants in the family *Gramineae* are a particularly preferred target plants for the methods of the invention.

Additional preferred targets include other commercially important crops, e.g., from the families *Compositae* (the largest family of vascular plants, including at

least 1,000 genera, including important commercial crops such as sunflower), and *Leguminosae* or the "pea family," which includes several hundred genera, including many commercially valuable crops such as pea, beans, lentil, peanut, yam bean, cowpeas, velvet beans, soybean, clover, alfalfa, lupine, vetch, *Lotus*, sweet clover, wisteria, and sweetpea.

Plants in the genera *Fragaria*, *Lotus*, *Medicago*, *Onobrychis*, *Trifolium*, *Trigonella*, *Vigna*, *Citrus*, *Linum*, *Geranium*, *Manihot*, *Daucus*, *Brassica*, *Raphanus*, *Sinapis*, *Atropa*, *Capsicum*, *Datura*, *Hyoscyamus*, *Lycopersicon*, *Nicotiana*, *Solanum*, *Petunia*, *Digitalis*, *Majorana*, *Cichorium*, *Helianthus*, *Lactuca*, *Bromus*, *Asparagus*, *Antirrhinum*, *Heterocallis*, *Nemesia*, *Pelargonium*, *Panicum*, *Pennisetum*, *Ranunculus*, *Senecio*, *Salpiglossis*, *Cucumis*, *Browaalia*, *Glycine*, *Lolium*, *Zea*, *Triticum*, *Sorghum*, *Malus*, and *Apium* are targets for mycotoxin detoxification, as are plants in the genera *Agrostis*, *Phleum*, *Dactylis*, *Sorgum*, *Setaria*, *Zea*, *Oryza*, *Triticum*, *Secale*, *Avena*, *Hordeum*, *Saccharum*, *Poa*, *Festuca*, *Stenotaphrum*, *Cynodon*, *Coix*, *Olyraeae*, *Phareae*, *Glycine*, *Pisum*, *Cicer*, *Phaseolus*, *Lens*, and *Arachis* are targets for acquiring mycotoxin detoxification activity.

Common crop plants which are targets for mycotoxin detoxification include corn, rice, triticale, rye, cotton, soybean, sorghum, wheat, oats, barley, millet, sunflower, canola, peas, beans, lentils, peanuts, yam beans, cowpeas, velvet beans, clover, alfalfa, lupine, vetch, *Lotus*, sweet clover, wisteria, sweetpea and nut plants (e.g., walnut, pecan, etc).

In construction of recombinant expression cassettes of the invention, a plant promoter fragment is optionally employed which directs expression of a nucleic acid in any or all tissues of a regenerated plant. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S transcription initiation region, the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens*, and other transcription initiation regions from various plant genes known to those of skill. Alternatively, the plant promoter may direct expression of the polynucleotide of the invention in a specific tissue (tissue-specific promoters) or may be otherwise under more precise environmental control (inducible promoters). Examples of tissue-specific promoters under developmental control include promoters that initiate transcription only

in certain tissues, such as fruit, seeds, or flowers.

Any of a number of promoters which direct transcription in plant cells can be suitable. The promoter can be either constitutive or inducible. In addition to the promoters noted above, promoters of bacterial origin which operate in plants include the octopine synthase promoter, the nopaline synthase promoter and other promoters derived from native Ti plasmids. See, Herrera-Estrella *et al.* (1983), *Nature*, 303:209-213. Viral promoters include the 35S and 19S RNA promoters of cauliflower mosaic virus. See, Odell *et al.* (1985) *Nature*, 313:810-812. Other plant promoters include the ribulose-1,3-bisphosphate carboxylase small subunit promoter and the phaseolin promoter. The promoter sequence from the E8 gene and other genes may also be used. The isolation and sequence of the E8 promoter is described in detail in Deikman and Fischer, (1988) *EMBO J.* 7:3315- 3327. Many other promoters are in current use and can be coupled to a mycotoxin detoxification nucleic acid to direct expression of the nucleic acid.

If polypeptide expression is desired, a polyadenylation region at the 3'-end of the coding region is typically included. The polyadenylation region can be derived from the natural gene, from a variety of other plant genes, or from, e.g., T-DNA.

The vector comprising the sequences (*e.g.*, promoters or coding regions) from genes encoding expression products of the invention will typically comprise a nucleic acid subsequence which confers a selectable phenotype on plant cells. The vector comprising the sequence will typically comprise a marker gene which confers a selectable phenotype on plant cells. For example, the marker may encode biocide tolerance, particularly antibiotic tolerance, such as tolerance to kanamycin, G418, bleomycin, hygromycin, or herbicide tolerance, such as tolerance to chlorosulfuron, or phosphinothricin (the active ingredient in the herbicides bialaphos and Basta). For example, crop selectivity to specific herbicides can be conferred by engineering genes into crops which encode appropriate herbicide metabolizing enzymes from other organisms, such as microbes. See, Padgett *et al.* (1996) "New weed control opportunities: Development of soybeans with a Round UP Ready™ gene" In: Herbicide-Resistant Crops (Duke, ed.), pp 53-84, CRC Lewis Publishers, Boca Raton ("Padgett, 1996"); and Vasil (1996) "Phosphinothricin-resistant crops" In: Herbicide-Resistant

Crops (Duke, ed.), pp 85-91, CRC Lewis Publishers, Boca Raton) (Vasil, 1996).

Transgenic plants have been engineered to express a variety of herbicide tolerance/metabolizing genes, from a variety of organisms-- and the expression of these genes in concert with the mycotoxin detoxification nucleic acid can be used a selectable marker for the presence of a vector comprising the mycotoxin detoxification nucleic acid. For example, acetohydroxy acid synthase, which has been found to make plants which express this enzyme resistant to multiple types of herbicides, has been cloned into a variety of plants (*see, e.g.,* Hattori, J., et al. (1995) Mol. Gen. Genet. 246(4):419). Other genes that confer tolerance to herbicides include: a gene encoding a chimeric protein of rat cytochrome P4507A1 and yeast NADPH-cytochrome P450 oxidoreductase (Shiota, et al. (1994) Plant Physiol. 106(1):17, genes for glutathione reductase and superoxide dismutase (Aono, et al. (1995) Plant Cell Physiol. 36(8):1687, and genes for various phosphotransferases (Datta, et al. (1992) Plant Mol. Biol. 20(4):619. Similarly, crop selectivity can be conferred by altering the gene coding for an herbicide target site so that the altered protein is no longer inhibited by the herbicide (Padgett, 1996). Several such crops have been engineered with specific microbial enzymes for confer selectivity to specific herbicides (Vasil, 1996). A wide variety of expression cassettes are known and available.

One of skill will recognize that after the expression cassette is stably incorporated in transgenic plants and confirmed to be operable, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

Computer System Elements

Software elements for manipulating strings of characters which correspond to mycotoxin detoxification nucleic acids can be used to direct synthesis of oligonucleotides relevant to shuffling of mycotoxin detoxification nucleic acids. Integrated systems comprising these and other useful features, e.g., one or more of: a digital computer with additional features such as high-throughput liquid control software, image analysis software, data interpretation software, a robotic liquid control armature for transferring solutions from a source to a destination (e.g., for manipulating selection assay solutions) operably linked to the digital computer, an input device (e.g., a computer keyboard) for

5 entering data to the digital computer to control high throughput liquid transfer by the
robotic liquid control armature an image scanner for digitizing label signals from labeled
10 assay components, or the like are a feature of the invention.

15 In one aspect, the invention provides an integrated system comprising a
5 computer or computer readable medium comprising a database having at least two artificial
homologous mycotoxin detoxification nucleic acid sequence strings, and a user interface
allowing a user to selectively view one or more sequence strings in the database. There are
15 a variety of sequence database programs for aligning and manipulating sequences. In
addition, standard text manipulation software such as word processing software (e.g.,
10 Microsoft Word™ or Corel WordPerfect™) and database software (e.g., spreadsheet
20 software such as Microsoft Excel™, Corel Quattro Pro™, or database programs such as
Microsoft Access™ or Paradox™) can be used in conjunction with a user interface (e.g., a
GUI in a standard operating system such as a Windows, Macintosh or LINUX system) to
25 manipulate strings of characters. In addition specialized alignment programs such as
15 BLAST can also be incorporated into the systems of the invention for alignment of
mycotoxin detoxification nucleic acids (or corresponding character strings).

30 In addition to the integrated system elements mentioned above, the
integrated system can also include an automated oligonucleotide synthesizer operably
linked to the computer or computer readable medium. Typically, the synthesizer is
20 programmed to synthesize one or more oligonucleotide comprising one or more
subsequence of one or more of the at least two artificial homologous mycotoxin
35 detoxification nucleic acids, e.g., as is useful in oligonucleotide shuffling procedures.

40 Modifications can be made to the method and materials as hereinbefore
described without departing from the spirit or scope of the invention as claimed, and the
25 invention can be put to a number of different uses, including:

The use of an integrated system to test monooxygenase in shuffled DNAs,
including in an iterative process.

45 An assay, kit or system utilizing a use of any one of the selection
strategies, materials, components, methods or substrates hereinbefore described. Kits
30 will optionally additionally comprise instructions for performing methods or assays,
packaging materials, one or more containers which contain assay, device or system
50 components, or the like.

5

10

5

15

In an additional aspect, the present invention provides kits embodying the methods and apparatus herein. Kits of the invention optionally comprise one or more of the following: (1) a shuffled component as described herein; (2) instructions for practicing the methods described herein, and/or for operating the selection procedure herein; (3) one or more mycotoxin assay component; (4) a container for holding mycotoxin detoxification nucleic acids or enzymes, other nucleic acids, transgenic plants, animals, cells, or the like and, (5) packaging materials.

20

10

In a further aspect, the present invention provides for the use of any component or kit herein, for the practice of any method or assay herein, and/or for the use of any apparatus or kit to practice any assay or method herein.

25

15

Accordingly, the disclosures and descriptions herein are intended to be illustrative, but not limiting, of the scope of the invention which is set forth in the following claims. One of skill will recognize many modifications which fall within the scope of the following claims. For example, all of the methods and compositions herein may be used in different combinations to achieve results selected by one of skill. *All publications and patent applications cited herein are incorporated by reference in their entirety for all purposes, as if each were specifically indicated to be incorporated by reference.*

30

35

40

45

50

55

Claims

5

10

15

20

25

30

35

40

45

50

55

5

WHAT IS CLAIMED IS:

10

15

20

25

30

35

40

45

50

55

- 1 1. A method of making a nucleic acid encoding a mycotoxin
2 detoxification activity, the method comprising:
3 recombining a plurality of parental nucleic acids to produce one or more
4 recombinant mycotoxin detoxification nucleic acid comprising a distinct or improved
5 mycotoxin detoxification activity; and,
6 selecting the one or more recombinant mycotoxin detoxification nucleic acid for
7 one or more encoded mycotoxin detoxification activity or,
8 selecting the one or more recombinant mycotoxin detoxification nucleic acid for
9 enhanced or reduced encoded polypeptide expression or stability;
10 thereby producing a selected shuffled mycotoxin detoxification nucleic acid,
11 which nucleic acid encodes a selected mycotoxin detoxification activity.
- 1 2. The method of claim 1, wherein the mycotoxin detoxification activity is
2 selected from: inactivation or modification of a polyketide, inactivation or modification of
3 an aflatoxin, inactivation or modification of a sterigmatocystin, inactivation or
4 modification of a trichothecene, and inactivation or modification of a fumonisin.
- 1 3. The method of claim 1, wherein the one or more recombinant
2 mycotoxin detoxification nucleic acid is selected by detecting a change in a physical
3 property of one or more mycotoxin in the presence of a polypeptide encoded by the one
4 or more mycotoxin detoxification nucleic acid, or by detecting cell growth or survival for a
5 cell transduced with the one or more mycotoxin detoxification nucleic acid, which cell is
6 cultured in the presence of the one or more mycotoxin.
- 1 4. The method of claim 1, wherein at least one of the parental nucleic
2 acids is the same as, or homologous to, a nucleic acid selected from: a monooxygenase, a
3 P450, trichothecene-3-O-acetyltransferase, a 3-O-Methyltransferase, a glutathione S-
4 transferase, an epoxide hydrolase, an isomerase, a macrolide-O-acyltransferase, a 3-O-
5 acetyltransferase, and a cis-diol producing monooxygenase which is specific for furan.

5

1 5. The method of claim 1, wherein the parental nucleic acids are
2 homologous.

10

1 6. The method of claim 1, wherein at least one of the parental nucleic
2 acids does not encode an anti-mycotoxin activity.

15

3 7. The method of claim 1, wherein the parental nucleic acid encodes a
4 polypeptide or polypeptide subsequence selected from: a monooxygenase, a P450,
5 trichothecene-3-O-acetyltransferase, a 3-O-Methyltransferase, a glutathione S-transferase,
6 an epoxide hydrolase, an isomerase, a macrolide-O-acetyltransferase, a 3-O-
7 acetyltransferase, and a cis-diol producing monooxygenase which is specific for furan.

20

25

1 8. The method of claim 1, wherein any of: the parental nucleic acids, the
2 one or more recombinant monooxygenase nucleic acid, and the selected recombinant
3 mycotoxin detoxification nucleic acid, is cloned into an expression vector.

30

1 9. A nucleic acid encoding a mycotoxin detoxification activity made by
2 the method of claim 1.

35

1 10. The method of claim 1, wherein the plurality of parental nucleic acids
2 are shuffled to produce a library of recombinant nucleic acids comprising one or more
3 library member nucleic acid encoding one or more mycotoxin detoxification activity,
4 which library is selected for one or more mycotoxin detoxification activity selected from:
5 inactivation or modification of a polyketide, inactivation or modification of an aflatoxin,
6 inactivation or modification of a sterigmatocystin, inactivation or modification of a
7 trichothecene, and inactivation or modification of a fumonisin, an increased ability to
8 chemically modify a mycotoxin, an increase in the range of mycotoxin substrates which a
9 polypeptide encoded by the nucleic acid can modify, an increased expression level of a
10 polypeptide encoded by the nucleic acid, a decrease in susceptibility of a polypeptide
11 encoded by the nucleic acid to protease cleavage, a decrease in susceptibility of a
12 polypeptide encoded by the nucleic acid to high or low pH levels, a decrease in

40

45

50

55

5
13 susceptibility of the protein encoded by the nucleic acid to high or low temperatures, and
14 a decrease in toxicity to a host cell of a polypeptide encoded by the selected nucleic acid.

10
1 11. A library of recombinant nucleic acids comprising one or more
2 monooxygenase activity made by the method of claim 10.

15
1 12. The library of claim 10, wherein the library is a phage display
2 library.

1 13. The method of claim 1, wherein the parental nucleic acids are
20 2 shuffled in a plurality of cells, which cells are prokaryotes or eukaryotes.

1 14. The method of claim 1, wherein the parental nucleic acids are
25 2 shuffled in a plurality of cells, which cells are plants, yeast, bacteria, or fungi.

1 15. The method of claim 1, wherein the parental nucleic acids are
2 shuffled in a plurality of cells; the method optionally further comprising one or more of:
30 3 (a) recombining DNA from the plurality of cells that display mycotoxin
4 detoxification activity with a library of DNA fragments, at least one of which undergoes
5 recombination with a segment in a cellular DNA present in the cells to produce
35 6 recombined cells, or recombining DNA between the plurality of cells that display
7 mycotoxin detoxification activity to produce cells with modified mycotoxin detoxification
8 activity;
9 (b) recombining and screening the recombined or modified cells to produce
40 10 further recombined cells that have evolved additionally modified mycotoxin detoxification
11 activity; and,
12 (c) repeating (a) or (b) until the further recombined cells have acquired a desired
45 13 mycotoxin detoxification activity.

1 16. A further recombined cell having acquired a desired mycotoxin
50 2 detoxification activity made by the method of claim 15.

5

1 17. The method of claim 1, wherein the method further comprises:

10

2 (a) recombining at least one distinct or improved recombinant nucleic acid with a
3 further mycotoxin detoxification activity nucleic acid, which further nucleic acid is the
4 same or different from one or more of the plurality of parental nucleic acids to produce a
5 library of recombinant mycotoxin detoxification nucleic acids;

15

6 (b) screening the library to identify at least one further distinct or improved
7 recombinant mycotoxin detoxification nucleic acid that exhibits a further improvement or
8 distinct property compared to the plurality of parental nucleic acids; and, optionally,

20

9 (c) repeating (a) and (b) until the resulting further distinct or improved
10 recombinant nucleic acid shows an additionally distinct or improved mycotoxin
11 detoxification property.

25

1 18. The method of claim 1, wherein the one or more recombinant
2 mycotoxin detoxification nucleic acid is present in one or more bacterial, yeast, plant or
3 fungal cells and the method comprises:

30

4 pooling multiple separate mycotoxin detoxification nucleic acids;
5 screening the resulting pooled mycotoxin detoxification nucleic acids to identify
6 distinct or improved recombinant mycotoxin detoxification nucleic acids that exhibit
7 distinct or improved mycotoxin resistance activity compared to a non-recombinant
8 mycotoxin resistance activity nucleic acid; and,
9 cloning the distinct or improved recombinant nucleic acid.

35

40

1 19. The method of claim 1, further comprising transducing the distinct or
2 improved nucleic acid into a prokaryote or eukaryote.

45

1 20. The method of claim 1, wherein recombining the plurality of parental
2 nucleic acids is performed by family gene shuffling.

50

1 21. The method of claim 1, wherein recombining the plurality of parental
2 nucleic acids comprises individual gene shuffling.

55

5

10

15

20

25

30

35

40

45

50

55

1 22. A plant transduced with the mycotoxin detoxification nucleic acid of
2 claim 22.

1 23. The plant of claim 22, wherein the plant is selected from the families
2 *Gramineae*, *Composite*, and *Leguminosae*.

1 24. The plant of claim 22, wherein the plant is selected from the genera:
2 *Fragaria*, *Lotus*, *Medicago*, *Onobrychis*, *Trifolium*, *Trigonella*, *Vigna*, *Citrus*, *Linum*,
3 *Geranium*, *Manihot*, *Daucus*, *Brassica*, *Raphanus*, *Sinapis*, *Atropa*, *Capsicum*, *Datura*,
4 *Hyoscyamus*, *Lycopersicon*, *Nicotiana*, *Solanum*, *Petunia*, *Digitalis*, *Majorana*,
5 *Cichorium*, *Helianthus*, *Lactuca*, *Bromus*, *Asparagus*, *Antirrhinum*, *Heterocallis*,
6 *Nemesia*, *Pelargonium*, *Panicum*, *Pennisetum*, *Ranunculus*, *Senecio*, *Salpiglossis*,
7 *Cucumis*, *Browaalia*, *Glycine*, *Lolium*, *Zea*, *Triticum*, *Sorghum*, *Malus*, *Apium*, *Agrostis*,
8 *Phleum*, *Dactylis*, *Sorgum*, *Setaria*, *Zea*, *Oryza*, *Triticum*, *Secale*, *Avena*, *Hordeum*,
9 *Saccharum*, *Poa*, *Festuca*, *Stenotaphrum*, *Cynodon*, *Coix*, *Olyrae*, *Phareae*, *Glycine*,
10 *Pisum*, *Cicer*, *Phaseolus*, *Lens*, and *Arachis*.

1 25. The plant of claim 22, wherein the plant is selected from corn, rice,
2 triticale, rye, cotton, soybean, sorghum, wheat, oats, barley, millet, sunflower, canola,
3 peas, beans, lentils, peanuts, yam beans, cowpeas, velvet beans, clover, alfalfa, lupine,
4 vetch, lotus, sweet clover, wisteria, sweetpea and a nut plant.

1 26. The plant of claim 22, wherein the plant exhibits mycotoxin
2 detoxification.

1 27. A DNA shuffling mixture, comprising: at least three homologous
2 DNAs, each of which is derived from a nucleic acid encoding a polypeptide or
3 polypeptide fragment which encodes mycotoxin detoxification activity.

1 28. The DNA shuffling mixture of claim 27, wherein the at least three

5

2 homologous DNAs are present in cell culture or *in vitro*.

10

1 29. A method of increasing mycotoxin detoxification of a cell,
2 comprising: performing whole genome shuffling of a plurality of genomic nucleic acids
3 in the cell and selecting for one or more mycotoxin detoxification activity.

15

1 30. The method of claim 29, wherein the genomic nucleic acids are from
2 a species or strain different from the cell.

20

1 31. The method of claim 29, wherein the cell is of prokaryotic or
2 eukaryotic origin.

25

30

35

40

45

50

55

1 32. The method of claim 29, wherein the mycotoxin detoxification
2 activity to be selected is selected from: inactivation or modification of a polyketide,
3 inactivation or modification of an aflatoxin, inactivation or modification of a
4 sterigmatocystin, inactivation or modification of a trichothecene, and inactivation or
5 modification of a fumonisin, an increased ability to chemically modify a mycotoxin, an
6 increase in the range of mycotoxin substrates for the cell, an increased expression level of
7 a mycotoxin detoxification polypeptide in the cell, a decrease in susceptibility of a
8 mycotoxin detoxification polypeptide in the cell to protease cleavage, a decrease in
9 susceptibility of a mycotoxin detoxification polypeptide in the cell to high or low pH
10 levels, a decrease in susceptibility of the protein in the cell to high or low temperatures,
11 and a decrease in toxicity to the cell by a mycotoxin detoxification polypeptide expressed
12 in the cell.